

Research Article

COEXISTENCE OF POLYMORPHISM IN FECUNDITY GENES BMPR1B AND GDF9 OF INDIAN KENDRAPADA SHEEP

Satyabrata Dash¹, Apratim Maity^{2*}, Purna Chandra Bisoi¹, Tapan Kumar Palai¹, Shamik Polley³,
Ayan Mukherjee⁴, Sachinandan De⁴

Received 27 February 2017, revised 20 April 2017

ABSTRACT: Present study was carried out to find out the status of mutations in three fecundity genes *i.e.* growth differentiation factor 9 (*GDF9/FecG*), bone morphogenetic protein 15 (*BMP15/FecX*) and bone morphogenetic protein receptor (*BMPR1B/FecB*) in Kendrapada sheep, the second most prolific sheep breed of India after Garole. Kendrapada ewes (n=85) were genotyped by Tetra-primer amplification refractory mutation system-PCR and a total of eleven SNP points over these three candidate fecundity genes (one point on *FecB* and five points each on *BMP15* and *GDF9*) were assessed. Out of eleven, two SNP points, *viz.* *FecB* and G4 of *GDF9* were found to be polymorphic in this breed. In this sheep breed average litter size of the ewes with non-carriers, heterozygous carrier and homozygous carrier of *FecB* locus mutation were 1.61, 1.80 and 2.06 respectively. G4 point of the *GDF9* gene was also polymorphic with average litter size of non-carriers, heterozygous carrier and homozygous carrier ewes were 1.63, 2.00 and 1.91 respectively. This study establishes Kendrapada sheep as the sixth sheep breed after Belclare/Cambridge, Lacaune, Small-tailed Han, Garole and Bayanbulak sheep, where coexisting polymorphism has been found in two different fecundity genes (*BMPR1B* and *GDF9* genes).

Key words: Prolificacy, *FecB*, *FecG*, *FecX*, T-ARMS PCR, Kendrapada sheep, India.

INTRODUCTION

In domesticated sheep (*Ovis aries*), ovulation rate and litter size vary widely among different breeds and within breeds. Although identifying the specific gene(s) behind this complex phenomenon was a formidable task, studies showed that a set of different genes, collectively called fecundity (*Fec*) genes (Piper and Bindon 1982, Davis *et al.* 1982, Davis 2005) play crucial role in determining ovulation rate and litter size of domesticated sheep. Three major *Fec* genes, having profound effect on the fecundity in sheep are bone morphogenetic protein receptor type IB (*BMPR1B*) or activin-like kinase 6 or *FecB* or Booroola gene (Souza *et al.* 2001), growth differentiation factor 9 (*GDF9*) or *FecG* (Hanrahan *et al.* 2004), and bone morphogenetic protein 15 (*BMP15*) or *FecX*

(Galloway *et al.* 2000). They are the member of transforming growth factor β (*TGF β*) super family (Juengel *et al.* 2004, Liu *et al.* 2014).

The *BMPR1B* gene is present in ovine chromosome 6 and it induces precocious maturation of ovarian follicles that ovulate at smaller size (McNatty *et al.* 1986). A G→A transition at nucleotide position 746 of *BMPR1B* cDNA produced 'hyperprolific' Booroola sheep, known as *FecB* (Souza *et al.* 2001, Wilson *et al.* 2001). The mutation in *FecB* allele is associated with the additive effect on ovulation rate and increase in litter size (Davis 2004, Kumar *et al.* 2006).

Bone morphogenic protein (*BMP15*) monitors a variety of cellular phenomena including oocyte

¹Department of Veterinary Biochemistry, College of Veterinary Science and Animal Husbandry, Orissa University of Agriculture and Technology, Bhubaneswar, Odisha 751003, India.

²Department of Veterinary Biochemistry, WBUAFS, Kolkata 700037, India.

³College of Veterinary Science and Animal Husbandry, Central Agricultural University, Jalukie, Peren, Nagaland 797110, India.

⁴Animal Genomics Lab., National Dairy Research Institute, Karnal, Haryana, India.

*Corresponding author. e - mail: toapu77@gmail.com

Table 1. Allelic and genotype frequencies of the Kendrapara sheep for *BMPR 1B* locus (FecB) and G4 locus of *GDF- 9* gene.

Gene (Mutation)	No. of animal	Allele frequency		Genotype frequency		
		G	A	GG	AG	AA
BMPR-1B (FecB)	85	0.60	0.40	0.353	0.494	0.153
GDF-9 (G4)	85	0.67	0.33	0.58	0.20	0.22

maturation and development (Galloway *et al.* 2002, Shimasaki *et al.* 2004). The gene plays crucial role in regulation of different granulosa cell processes and folliculogenesis. Mutations in the mature *BMP15* coding region have been described (Davis 2005) with a positive impact on prolificacy of various sheep breeds (Hanrahan *et al.* 2004). One copy of the Inverdale (*FecX^I*), Hanna (*FecX^H*), Belclare (*FecX^B*), Galway (*FecX^G*) or Lacaune (*FecX^L*) allele cause increase in about 0.6 lambs per ewe, compared to the wild type (Hanrahan *et al.* 2004, Bodin *et al.* 2007).

Ovine Growth differentiation factor 9 (*GDF9*), a close autosomal paralogue of *BMP15*, has been mapped to sheep chromosome 5 (Sadighi *et al.* 2002). *GDF9* and *BMP15* are supposed to form non-covalent homo and heterodimers *in vivo* and modulate the ovulation rate in species-specific way (Moore *et al.* 2004). Throughout the coding region of the *GDF9* gene eight single nucleotide polymorphisms (G1-G8) have been identified (Hanrahan *et al.* 2004). Out of these eight mutation points three points (G2, G3, and G5) are single nucleotide changes that are synonymous in nature. A recent study on *GDF9* gene in Garole sheep revealed that 85.71 percent of studied animals had AA genotype while only 14.29 percent animals had AB genotype (Raja *et al.* 2016).

The *BMPR1B*, together with *BMPRII*, execute *BMP15* signalling in the ovarian follicles (Moore *et al.* 2003). *GDF9* and *BMP15* have functional significance during cumulus expansion, oocyte maturation and ovulation (Elvin *et al.* 1999, Gui and Joyce 2005, Yoshino *et al.* 2006). Besides these, two paracrine factors control follicle growth (Dong *et al.* 1996, Nilsson and Skinner 2002), cumulus and granulosa cell proliferation (Hayashi *et al.* 1999, Gilchrist *et al.* 2006, Spicer *et al.* 2006), cell-survival signalling (Hussein *et al.* 2005, Orisaka *et al.* 2006) and act as regulator of numerous growth factors and endocrine hormones (Juengel *et al.* 2004). So study to find out existence of polymorphisms in these three important fecundity related genes would be useful to increase the reproductive efficiency of animal through genetic manipulation.

Kendrapada sheep is a prolific breed with its natural abode in coastal districts like Kendrapada, Jagatsinghpur

and Cuttack in Odisha, India. Majority of pure bred animals are found in Kendrapada district. The family flock size varies greatly, from as low as 5 to as high as 50. Males and females are maintained in ratio of 1:30 to 1:40. The age of sexual maturity is about 349 days and age at first lambing is about 518 days. The sheep breeds throughout the year but most of them come to heat in summer. Kendrapada sheep produces 35.1% single, 62.6% twin and 2.3% triplet (Patro *et al.* 2006). Recent study by Kumar *et al.* (2008) confirmed presence of FecB mutation in the ewes with increased prolificacy of this breed. It would be interesting to find out whether increased prolificacy of this breed is associated with mutations in other fecundity candidate genes like *BMP15* and *GDF9*, along with FecB.

The objectives of the current study were to detect the polymorphism of *GDF9*, *BMP15* and *BMPR1B* genes in Kendrapada sheep and to investigate their association with prolificacy of the breed.

MATERIALS AND METHODS

A total of 85 Kendrapada ewes, with a history of multiple births (twins and triplets), were identified from different villages of Kendrapada district, Odisha for our study. About 10 ml of venous jugular blood sample was collected from each animal in sterile vacutainer containing K₃-EDTA as an anticoagulant. The collected blood samples were stored at 4°C and transported to the laboratory at the earliest. After processing, the genomic DNA was isolated from white blood cells using standard phenol-chloroform procedure (Sambrook and Russell 2001). The DNA samples were dissolved in TE buffer (pH 8.0) and stored at -20°C before use.

Tetra-primer amplification refractory mutation system polymerase chain reaction (T-ARMS-PCR), as described by Ye *et al.* (2001), was carried out to detect the FecB point mutation of *BMPR1B* gene. Briefly two pairs of primers were designed to amplify fragments of differing sizes, each representing one allele. Amplicons were easily resolved by standard agarose gel electrophoresis. A total of ten point mutations of other two important candidate genes, five each in *GDF9* (G1, G4, G6, G7 and G8) and *BMP15* (*FecX^G*, *FecX^H*, *FecX^I*, *FecX^L* and *FecX^B*) gene

Table 2. Effect of different genotype of FecB and G4 on lambing size.

Gene	SNP	Genotype	No. of animals	Average litter size
B M P R- IB	Fec B	AA	13	1.61
		AG	42	1.80
		GG	30	2.06
G D F 9	G 4	AA	19	1.63
		AG	17	2.00
		GG	49	1.92

were screened by T-ARMS-PCR as mentioned above using the primers described earlier (Polley *et al.* 2009, 2010).

PCR reaction was performed in a 25 µl reaction mixture containing 2.5µl of 10× Paq5000 reaction buffer (provides a final Mg²⁺ concentration of 2mM), 200 µM of dNTPs (Fermentas, Lithuania), 5 pmol of each primer and 0.5 units of Paq5000™ DNA polymerase (Stratagene, USA). About 50 ng of genomic DNA was used as template. The PCR cycling parameters were optimized separately for detecting each of the allele specific amplifications. The PCR products were separated by horizontal submarine agarose gel (2-3%, free from DNase and RNase, Sigma, USA) electrophoresis in 1× TAE buffer at 100 V. The gel was stained with ethidium bromide solution (0.5µg/ml), maintained for 10 min. in darkness and photographed using a molecular imager (Gel Doc XR, BIORAD).

RESULTS AND DISCUSSION

The molecular crosstalk among BMP15, BMPR1B and GDF9 and their liaison with female fecundity in domesticated sheep is a well-acquainted fact. Recently, G to A nucleotide substitution was found in GDF9 locus that affects litter size in Iranian Baluchi sheep (Moradband *et al.* 2011). Two mutations including *BMPR-IB/FecB* and *GDF9/G1* were also found in Bayanbulak sheep (Zuo *et al.* 2013).

BMPR1B: In BMPR1B gene, 'A' allele (non-carrier) represents the wild type and 'G' allele (carrier) represents the mutant type (FecB). Genomic DNA samples from 85 Kendrapada sheep were analyzed and all three possible genotypes, AA, AG and GG were observed. These three genotypes (AA, AG and GG) represent the wild homozygous, mutant heterozygous and mutant homozygous respectively. Presence of each genotype was confirmed by visualizing amplicon size variation following PCR. Amplicon sizes of 1178 bp (common outer) and 1097 bp ('A' allele specific) confirmed AA genotype. In case of mutant carrier 'GG' sheep 1178bp




Fig. 1. Agarose gel electrophoresis (2%) of tetra-primer amplification refractory mutation system-PCR amplicons for FecB locus of BMPR1B gene.

The size of common outer product (1178 bp) and allele-specific wild-type (1097 bp) and mutant-type (136 bp) inner product are indicated by arrow on the left side of the gel photograph. Lane L: DNA molecular weight marker (O'RangeRular 100bp DNA Ladder, Fermentas, Lithuania) indicated by arrow on the right side of the gel photograph.




Fig. 2. Agarose gel electrophoresis (2.5%) of tetra-primer amplification refractory mutation system-PCR amplicons for G4 locus of GDF9 gene.

The size of common outer product (417 bp) and allele-specific wild-type (261 bp) and mutant-type (212 bp) inner product are indicated by arrow on the right side of the gel photograph. Lane L: DNA molecular weight marker (O'RangeRular 100bp DNA Ladder, Fermentas, Lithuania) indicated by arrow on the left side of the gel photograph.

and 136 bp ('G' allele specific) products were amplified. However, in heterozygous sheep 1178 bp, 1097 bp and 136 bp products were amplified (Fig. 1). The overall genotype frequencies of AA, AG and GG were found to be 0.153, 0.494 and 0.353 respectively. The allelic frequencies for A and G alleles were 0.40 and 0.60

respectively (Table 1). The results indicate that about 85% of the individuals in the study population are carrying mutation. Frequency of the mutant type allele is more than the wild type allele frequency (0.6 vs. 0.4) but is less than 0.73, determined earlier by Kumar *et al.* (2008) in the same breed in smaller sized flock. Average litter sizes of non-carrier wild types, heterozygous and mutant ewes were 1.61, 1.80 and 2.06 respectively (Table 2). The lambing records of the experimental sheep reveals that average litter size is the highest in homozygous mutant type followed by heterozygous mutant and homozygous wild type animals. This finding is in line with the other findings related to the phenotypic effect of FecB mutation in sheep. This is the sixth sheep breed in the world after Booroola, Garole, Javanese, Hu and small tailed Han carrying the particular point mutation that has impact on the litter size (Kumar *et al.* 2008).

BMP15: Earlier studies pointed out that the ewes with mutations in five different points (FecX^I, FecX^H, FecX^L, FecX^G and FecX^B) of BMP15 gene showed increased ovulation rate in heterozygous condition and sterility in homozygous condition due to impaired follicular development (Davis *et al.* 1982, Bodin *et al.* 2002, Hanrahan *et al.* 2004). In the current study, same five point mutations (Q239ProTer, Q23Ter, V31D, C53Y and S99I) in BMP15 were typed by tetra-primer ARMS-PCR in Kendrapada sheep. However, in our study, we did not find mutation in any of these five points. It implicates that increased litter size in breed has no association with the five known SNPs of BMP15. Several instances are there where no BMP15 mutations were found in these points in the different prolific sheep breeds of the world. None of Small Tailed Han and Hu sheep carried the FecX^B or FecX^H mutation (Liu *et al.* 2003, Chu *et al.* 2005^{a,b}). None of Garole, Javanese, Small Tailed Han, or Hu sheep had the Fec X^I mutation (Davis *et al.* 2002, Liu *et al.* 2003, Chu *et al.* 2005^a). The prolific Hu sheep did not have the FecX^G mutation (Chu *et al.* 2005^a). The present report is also in agreement with the recent finding of Kumar *et al.* (2008) that Kendrapada sheep with high litter size and twinning rate is not carrying FecX^G mutation.

GDF9: Only G4 point was found to be polymorphic in Kendrapada sheep after screening five point mutations in GDF9 gene at five positions (G1, G4, G6, G7 and G8) (Fig. 2). Hanrahan *et al.* (2004) reported that change in a specific point mutation in GDF9 gene (G8) causes an amino acid substitution that leads to increased prolificacy in Belclare and Cambridge sheep. Ewes with heterozygous GDF9 (G8) mutation have an increased ovulation rate but infertile in case of homozygous mutant (Juengel *et al.* 2004). In the present study, both 'A' and 'G' alleles were detected in G4 locus with frequencies of

0.33 and 0.67 respectively. All three possible genotypes were observed (AA, AG and GG) with genotype frequency of 0.22, 0.20 and 0.58 respectively (Table 1). The mutant (GG) genotype of G4 point was found to be the predominant genotype in tested Kendrapada population. Our results showed that ~80% of the individuals under study were carrying mutation in the G4 point of the gene. Average litter size of these three genotypes non-carriers wild types, heterozygous and mutant ewes had 1.63, 2.00 and 1.91 respectively (Table 2). Here is an interesting finding that presence of single copy of mutant allele in the G4 SNP point of GDF9 gene *i.e.*, in the heterozygous mutant animal shows maximum average litter size. Available literature suggests partly similar kind of observation in case of GDF 9 gene and conferred the phenomenon as a case of heterozygote advantage (Dong *et al.* 1996, Gemmell and Slate 2006). However, consolidated data collected over many years on ovulation rates, litter size and lambing rates of this local sheep breed is required to conclude the present case as compelling example of heterozygote advantage.

CONCLUSION

Finally, the results indicate that among 11 SNP points on three fecundity genes, at least two loci, FecB loci of BMPR1B gene and G4 of GDF9, are polymorphic in the Kendrapada sheep. Therefore, Kendrapada sheep is the sixth sheep breed after Belclare/Cambridge, Lacaune, Small-tailed Han, Garole and Bayanbulak where coexisting polymorphism has been found in two different fecundity genes. Further studies are needed to draw conclusion on the origin of these mutations and whether it is influenced by Garole or some other local breeds. Profitable and sustainable sheep production through genetic manipulation is a high priority for India. Reproductive performance of non-prolific local sheep breeds of India carrying other desired traits can be optimized through incorporation of prolificacy genes by advanced breeding strategy. Therefore, apart from Garole, Kendrapada may be the breed of choice for improving the prolificacy of Indian sheep breeds.

REFERENCES

- Bodin L, SanCristobal M, Lecerf F, Mulsant P, Bibe B, Lajous D, Belloc JP, Eychenne F, Amigues Y, Elsen JM (2002) Segregation of a major gene influencing ovulation in progeny of Lacaune meat sheep. *Genet Sel Evol* 34: 447- 464.
- Bodin L, Di Pasquale E, Fabre S, Bontoux M, Monget P, Ersani L, Mulsant P (2007) A novel mutation in the Bone Morphogenetic Protein 15 gene causing defective protein secretion is associated with both increased

- ovulation rate and sterility in Lacaune sheep. *Endocrinol* 148(1): 393-400.
- Chu MX, Cheng R, Chen GH, Fang L, Ye SC (2005a) Study on bone morphogenetic protein 15 as a candidate gene for prolificacy of Small Tailed Han sheep and Hu sheep. *J Anhui Agr Uni* 32: 278-282.
- Chu MX, Sang LH, Wang JY, Fang L, Ye SC (2005b) Study on BMP15 and GDF 9 as candidate genes for prolificacy of Small Tailed Han sheep. *Acta Genet Sin* 32: 38-45.
- Davis GH (2004) Fecundity genes in sheep. *Anim Reprod Sci* 82-83: 247-253.
- Davis GH (2005) Major genes affecting ovulation rate in sheep. *Genet Sel Evol* 37: S11-S23.
- Davis GH, Montgomery GW, Allison AJ, Kelly RW, Bray AR (1982) Segregation of a major gene influencing fecundity in progeny of Booroola sheep. *New Zealand J Agr Res* 25: 525-529.
- Davis GH, Galloway SM, Ross IK *et al.* (2002) DNA Tests in prolific sheep from eight countries provide new evidence on origin of the Booroola (FecB) mutation. *Biol Reprod* 66: 1869-1874.
- Dong J, Albertini DF, Nishimori K, Kumar TR, Lu N, Matzuk MM (1996) Growth differentiation factor-9 is required during early ovarian folliculogenesis. *Nature* 383: 531-535.
- Elvin JA, Clark AT, Wang P, Wolfman NM, Matzuk MM (1999) Paracrine actions of growth differentiation factor-9 in the mammalian ovary. *Mol Endocrinol* 13: 1035-1048.
- Galloway SM, Gregan SM, Wilson T, McNatty KP, Juengel JL, Ritvos O, Davis GH (2002) Bmp15 mutations and ovarian function. *Mol Cell Endocrinol* 191: 15-18.
- Gemmell NJ, Slate J (2006) Heterozygote advantage for fecundity. *PLoS ONE* 1(1): e125.
- Gilchrist RB, Ritter LJ, Myllymaa S, Kaivo-Oja N, Dragovic RA, Hickey TE, Ritvos O, Mottershead DG (2006) Molecular basis of oocyte-paracrine signalling that promotes granulosa cell proliferation. *J Cell Sci* 119: 3811-3821.
- Gui LM, Joyce IM (2005) RNA interference evidence that growth differentiation factor-9 mediates oocyte regulation of cumulus expansion in mice. *Biol Reprod* 72: 195-199.
- Hanrahan PJ, Gregan SM, Mulsant P, Mullen M, Davis GH, Powell R, Galloway SM (2004) Mutations in the genes for oocyte-derived growth factors GDF9 and BMP15 are associated with both increased ovulation rate and sterility in Cambridge and Belclare sheep (*Ovis aries*). *Biol Reprod* 70(4): 900-909.
- Hayashi M, McGee EA, Min G, Klein C, Rose UM, van Duin M, Hsueh AJ (1999) Recombinant growth differentiation factor-9 (GDF-9) enhances growth and differentiation of cultured early ovarian follicles. *Endocrinol* 140: 1236-1244.
- Hussein TS, Froiland DA, Amato F, Thompson JG, Gilchrist RB (2005) Oocytes prevent cumulus cell apoptosis by maintaining a morphogenic paracrine gradient of bone morphogenetic proteins. *J Cell Sci* 118: 5257-5268.
- Juengel JL, Hudson NL, Whiting L, McNatty KP (2004) Effects of immunization against bone morphogenetic protein 15 and growth differentiation factor 9 on ovulation rate, fertilization, and pregnancy in ewes. *Biol Reprod* 70: 557-561.
- Kumar S, Kolte AP, Mishra AK, Singh VK (2006) Evidence that twinning in Malpura sheep is not linked with Booroola fecundity (FecB) gene. *Ind J Anim Sci* 76(9): 764-767.
- Kumar S, Mishra AK, Kolte AP, Dash SK, Karim SA (2008) Screening for Booroola (FecB) and Galway (FecX^G) mutation in Indian sheep. *Small Rum Res* 80: 57-61.
- Liu SF, Jiang YL, Du LX (2003) Studies of BMPR-IB and BMP15 as candidate genes for fecundity in Little Tailed Han sheep. *Yi Chuan Xue Bao* 30(8): 755-760.
- Liu Q, Pan Z, Wang X, Hu W, Di R, Yao Y, Chu M (2014) Progress on major genes for high fecundity in ewes. *Front Agr Sci Eng* 1(4): 282-290.
- McNatty KP, Lun S, Heath DA, Ball K, Smith P, Hudson NL, McDiarmid J, Gibb M, Henderson KM (1986) Differences in ovarian activity between Booroola Merino ewes which were homozygous, heterozygous and non-carriers of a major gene influencing their ovulation rate. *J Reprod Fertil* 77: 193-205.
- Moore RK, Erickson GF, Shimasaki S (2004) Are BMP-15 and GDF-9 primary determinants of ovulation quota in mammals? *Trends Endocrinol Metab* 15: 356-361.
- Moore RK, Otsuka F, Shimasaki S (2003) Molecular basis of bone morphogenetic protein-15 signaling in granulosa cells. *J Biol Chem* 278: 304-310.

- Moradband F, Rahimi G and Gholizadeh M (2011) Association of Polymorphisms in Fecundity Genes of GDF9, BMP15 and BMP15-1B with Litter Size in Iranian Baluchi Sheep. *Asian-Aust J Anim Sci* 24(9): 1179-1183.
- Nilsson EE, Skinner MK (2002) Growth and differentiation factor-9 stimulates progression of early primary but not primordial rat ovarian follicle development. *Biol Reprod* 67: 1018-1024.
- Orisaka M, Orisaka S, Jiang JY, Craig J, Wang Y, Kotsuji F, Tsang BK (2006) Growth differentiation factor 9 is antiapoptotic during follicular development from preantral to early antral stage. *Mol Endocrinol* 20: 2456-2468.
- Patro BN, Mallick CR, Rao PK, Panda P (2006) Production performance of indigenous meat type sheep in Kendrapada district of coastal Orissa. *Ind J Small Rum* 12(1): 42-47.
- Piper LR, Bindon BM (1982) The Booroola Merino and the performance of medium non-peppin crosses at Armidale. *Proceedings of a Workshop. CSIRO, 24-25 August' 1980, Armidale.* 09-19.
- Polley S, De S, Batabyal S, Kaushik, R, Yadav P, Arora JS, Chattopadhyay S, Pan S, Brahma B, Datta TK, Goswami SL (2009) Polymorphism of fecundity genes (BMPR1B, BMP15 and GDF9) in the Indian prolific Black Bengal goat. *Small Rum Res* 85(2): 122-129.
- Polley S, De S, Brahma B, Mukherjee A, Vinesh PV, Batabyal S, Arora JS, Pan S, Samanta AK, Datta TK, Goswami SL (2010) Polymorphism of BMPR1B, BMP15 and GDF9 fecundity genes in prolific Garole sheep. *Trop Anim Health Prod* 42(5): 985-993.
- Raja D, Taraphder S, Sahoo AK, Paul R, Raj Kumar (2016) PCR-SSCP analysis of growth differentiation Factor-9 gene of Garole sheep. *Explor Anim Med Res* 6(1): 37-46.
- Sadighi M, Bodensteiner KJ, Beattie AE, Galloway SM (2002) Genetic mapping of ovine growth differentiation factor 9 (GDF9) to sheep chromosome 5. *Anim Genet* 33: 244-245.
- Sambrook J, Russell D (2001) *Molecular cloning: A laboratory manual.* Cold Spring Harbor Press, NY.
- Shimasaki S, Moore RK, Otsuka F, Erickson GF (2004) The bone morphogenetic protein system in mammalian reproduction. *Endocr Rev* 25: 72-101.
- Souza CJ, MacDougall C, Campbell BK, McNeilly AS, Baird DT (2001) The Booroola (FecB) phenotype is associated with a mutation in the bone morphogenetic receptor type 1 B (BMPR1B) gene. *J Endocrinol* 169: R1-R6.
- Spicer LJ, Aad PY, Allen D, Mazerbourg S, Hsueh AJ (2006) Growth differentiation factor-9 has divergent effects on proliferation and steroidogenesis of bovine granulosa cells. *J Endocrinol* 189: 329-339.
- Wilson T, Wu XY, Juengel JL, Ross IK, Lumsden JM, Lord EA, Dodds KG, Walling GA, McEwan JC, O'Connell AR, McNatty KP, Montgomery GW (2001) Highly prolific Booroola sheep have a mutation in the intracellular kinase domain of bone morphogenetic protein IB receptor (ALK-6) that is expressed in both oocytes and granulosa cells. *Biol Reprod* 64: 1225-1235.
- Ye S, Dhillon S, Ke X, Collins AR, Day IN (2001) An efficient procedure for genotyping single nucleotide polymorphisms. *Nucleic Acids Res* 29(17): E 88-89.
- Yoshino O, McMahon HE, Sharma S, Shimasaki S (2006) A unique preovulatory expression pattern plays a key role in the physiological functions of BMP-15 in the mouse. *P Natl Acad Sci* 103: 10678-10683.
- Zuo B, Qian H, Wang Z, Wang X, Nisa N, Bayier A, Ying S, Hu X, Gong C, Guo Z and Wang F (2013) A Study on *BMPR-IB* Genes of Bayanbulak Sheep. *Asian-Aust J Anim Sci* 26(1): 36-42.

***Cite this article as:** Dash S, Maity A, Bisoi PC, Palai TK, Polley S, Mukherjee A, De S (2017) Coexistence of polymorphism in fecundity genes BMPR1B and GDF9 of Indian Kendrapada sheep. *Explor Anim Med Res* 7(1): 33-38.