

Research Article

## ANTIDEPRESSANT PLANTS AND THEIR USE BY THE *CHIRU* TRIBES OF MANIPUR, INDIA

Soibam Puspa Chanu\*, Ranjana Rajkumari, Athokpam Pinokiyo

Received 02 January 2023, revised 06 August 2023

**ABSTRACT:** Depression is a potentially life-threatening mental disorder with serious socioeconomic consequences. It is very common and affects people of all age groups. Clinically, a class of synthetic drugs called antidepressants is used to treat depression but face limitations due to multiple side effects. Traditional healing practices followed by different communities worldwide use plants as a source of antidepressants. The Chiru tribe is one such community with rich ethnomedicinal knowledge having their unique ways to cure mental ailments. Ten plants [*Canthium parviflorum* Lam., *Centella asiatica* (L.) Urb., *Curcuma longa* L., *Holmskioldia sanguinea* Retz., *Magnolia champaca* (L.) Baill. ex Pierre, *Marsilea minuta* L., *Ocimum tenuiflorum* L., *Phyllanthus emblica* L., *Rotheca serrata* (L.) Steane & Mabb. and *Tagetes lucida* Cav.)] are used by Chiru tribes as potential antidepressants. Out of these, two plants viz., *Canthium parviflorum* Lam. and *Holmskioldia sanguinea* Retz. are recorded as antidepressants for the first time by the Chiru tribes. Two antidepressant plants (*Canthium parviflorum* Lam. and *Curcuma longa* L.) are used for treating memory loss. A total of 132 medicinal plants have been recorded in literature to contain phytochemicals with antidepressant activity worldwide. Changes in lifestyle and lack of properly documented data on medicinal plants pose a serious concern. The plant-based alternatives to treat depression will be much safer and cost-effective in the long run. The primitive knowledge could prove useful in developing potent antidepressants shortly. For this, further productive research and appropriate conservation methods for these plant species must be initiated.

**Key words:** Antidepressant plants, Chiru tribe, Ethnomedicinal knowledge, Phytochemicals.

### INTRODUCTION

Use of herbs as some medicine was started at very ancient stage of human civilization and validation of documented uses of such plant medicines is a very important area of present-day research performed for their possible effective use in different forms (Pattanayak 2021a, Patel *et al.* 2022, Paul and Sujata 2022). The ethnic communities and the people living in remote places use such plant resources as medicine and the collection of information from them is still very important (Pradhan *et al.* 2021). The present study was conducted on a tribe of Monipur, India to get information related to the use of plants as some medicine for anxiety, mental disorder, and alike problems.

Among mental disorders, depression is one of the most commonly known psychological disorders

affecting thoughts, feelings, behavior, and sense of well-being in a person (Pannu *et al.* 2021). With rising cases of mental illness and suicidal tendencies being part of it, it presents a significant socioeconomic burden (Pannu *et al.* 2021, Bondy *et al.* 2002) It is so common that it is expected to be the primitive cause of various other diseases by 2030, as per WHO report (Pannu *et al.* 2021). Clinically, synthetic drugs are used to treat depression but they have several limitations like multiple side effects, slow onset of action, poor remission and response rates, and drug-drug and dietary interactions (Bondy *et al.* 2002, Khushboo *et al.* 2017, Pardhe *et al.* 2020). Several emerging pieces of research have shown an association between dietary measures and the risk of depression pointing to the use of nutraceuticals as antidepressants (Pannu *et al.* 2021, Hritcu *et al.* 2017, Martins 2018).

---

Department of Botany, DM College of Science, Dhanamanjuri University, Imphal-795001, India.

\*Corresponding author. e-mail: chanupusasoibam@gmail.com

Several phytochemicals in plants have been studied for antidepressant properties. The Indian traditional system of medicine AYUSH explores a wide variety of medicinal and aromatic plants to treat various ailments with minimum side effects and low costs. The agroclimatic conditions of Manipur sustain the diversity of medicinal plants in the region. Since time immemorial, different ethnic groups inhabiting different regions of the state have followed traditional healthcare practices with their rich knowledge of medicinal plants. These age-old home remedies and associated medico-religious practices are often cheap, safe, long-time tested, and based on local resources and strength. The *Chiru* tribe of Manipur too has a repository of knowledge on medicinal plants, amongst which is their indigenous knowledge to cure mental illness. The WHO has even started documenting and listing popular tribal remedies used to cure ailments. The primitive knowledge could prove useful to discover chemically useful compounds which could be prototype compounds to treat psychological disorders including several forms of depression. There is a future need for pre-clinical and clinical trials to prove their efficacy as antidepressants and further development of therapeutically useful drugs.

The *Chiru* tribe is one of the *Kuki* Scheduled tribe groups of Manipur inhabiting 3 districts and 13 villages, viz. one in Churachandpur district (Charoi Khullen), two in Tamenglong district (Lamdangmei and Dolang) and ten in Senapati district (Dolang Khunou, Nugsai, Bungte Khullen, Lower Bungte, Sadu *Chiru*, Thangjing *Chiru*, Kangchup *Chiru*, Waithou, Uran *Chiru* and Chawang *Chiru*), of Manipur in different geographical regions. The traditional knowledge of *Chirus* is not documented properly yet. No proper investigation has been done so far to explore the traditional knowledge of antidepressants of the *Chiru* tribes. Traditional knowledge faces the risk of disappearing before documentation due to changes in livelihood patterns and environmental degradation. Hence, the present study has been carried out.

## METHODS

Five villages, viz., Bungte Khullen, Kangchup *Chiru*, Lower Bungte Khullen, Sadu *Chiru*, and Uran *Chiru* in the Senapati district of Manipur were chosen as the study area. A total of 20 field-cum survey tours were conducted throughout the five villages. The *Chiru* tribe is most populated in these villages. Prior Informed Consent (PIC) was taken from the headmen of the

villages and the villagers after explaining the purpose of the study. With regards to sharing of information on traditional Health Practices, an agreement was made with due compliance to the norms of the North Eastern Institute of Folk Medicine (NEIFM), Department of AYUSH. The informants were interviewed with queries including the description of the health condition, diagnostic features used by the healer, details of treatment followed, the local name of the plant, the plant part used, preparation of medicine and dosage (in terms of quantity and frequency of intake) recommended to cure the ailment. In this way, proper data was collected using Participatory Rural Appraisal (PRA) and interview-questionnaire methods (Martin 2007). Meetings were held in the presence of headmen and villagers to share the knowledge of the traditional healthcare practitioners.

With the help of the *Chiru* tribe, a direct visit was made to the nearby forest to identify and collect the medicinal plants used as antidepressants. Further, the collected plant species were identified with the help of regional floras in consultation with authentic herbarium at Botanical Survey of India, Kolkata and Botanical Survey of India, Eastern Regional Centre, Shillong, and herbarium at DM College of Science, Imphal, Manipur. Also, herbarium specimens were prepared and deposited in the herbarium at DM College of Science, Dhanamanjuri University Imphal, Manipur, India.

## RESULTS AND DISCUSSION

A literature review resulted in the documentation of around 130 antidepressant plants (Table 1). Altogether 10 plant species have been used by *Chiru* tribes for treating mental disorders, out of which two are recorded for the first time (Marked by \*).

The uses of the plants with their dosage are described below:

1. \**Canthium parviflorum* Lam. (Local name-*Lam-heibi*, Family- Rubiaceae):

When the leaf extract of the plant (dosage- 1 teaspoon) is given daily to the patient for about 2-3 months, it reduces loss of memory.

2. *Centella asiatica* (L.) Urb. (Local name- *Peruk*, Family- Apiaceae):

Consumption of fresh juice of the whole plant (mixed with water) for 2-3 weeks reduces mental disorders.

3. *Curcuma longa* L. (Local name- *Yaingang*, Family- Zingiberaceae):

## Antidepressant plants and their use by the Chiru Tribes of Manipur, India

When aqueous extract of the plant rhizome is administered orally 2 times daily for 2-3 weeks, it increases memory and reduces depression.

4. \**Holmskioldia sanguinea* Retz. (Local name- *Kharamleishok*, Family- Verbenaceae):

Boiled extract of the young shoots is used as an antidepressant.

5. *Magnolia champaca* (L.) Baill. ex Pierre (Local name- *Leihao*, Family- Magnoliaceae):

The bark of the tree with ginger rhizome when ground and mixed with water is administered to the patient for reducing mental disorder.

6. *Marsilea minuta* L. (Local name- *Ishing Yenshang*, Family- Marsileaceae):

Decoction of the whole plant mixed with tea is consumed for recovering from mental disorders.

7. *Ocimum tenuiflorum* L. (Local name- *Tulsi*, Family- Lamiaceae):

Boiled extract of the leaves and inflorescence of the plant is given daily for mental disorders until recovery.

8. *Phyllanthus emblica* L. (Local name- *Heigrü*, Family- Phyllanthaceae):

Fresh juice of the fruit mixed with water is used to treat psychiatric disorders.

9. *Rotheca serrata* (L.) Steane & Mabb. (Local name- *Moirang Khanamba*, Family-Verbenaceae)

Fresh root and leaf of the plant when crushed with water is used to reduce mental disorder.

10. *Tagetes lucida* Cav. (Local name- *Sanalei thondabi*, Family- Asteraceae):

Aqueous extract of the leaves is used as an antidepressant.

Out of these ten antidepressant plants, two plants (*Canthium parviflorum* Lam. and *Curcuma longa* L.) are used for treating memory loss. Out of the ten species of plants, eight plants, viz. *Centella asiatica* (L.) Urb., *Curcuma longa* L., *Magnolia champaca* (L.) Baill. Ex Pierre., *Marsilea minuta* L., *Ocimum tenuiflorum* L., *Phyllanthus emblica* L., *Rotheca serrata* (L.) Steane & Mabb. and *Tagetes lucida* Cav. are also reported to have antidepressant properties by earlier studies. Antidepressant plants, *Canthium parviflorum* Lam. (Rubiaceae) used to reduce loss of memory, and *Holmskioldia sanguinea* Retz. (Verbenaceae) used to reduce mental disorders are reported for the first time by the *Chiru* tribe.

The plant parts used for treatment included the whole plant, leaves, fruits, roots, rhizome, the bark of the tree, and inflorescence, leaves being mostly used. The *Chiru* tribes collected the medicinal plants from their immediate environment or nearby forests and prepared their traditional medicines afresh as and when required. They do not have any well-recognized mechanism to preserve and store their traditional medicines. They have strong confidence in their traditional medicines though they could not explain how the plants are effective in treating mental ailments. They also ascertain that the plant extracts in the recommended doses pose no established side effects throughout their age-old practice.

It is observed that in most cases, they use the plant parts directly after use, without any processing or modification. It is a very important aspect of the use of plants for medicinal purposes, as validation of reported uses of plant parts is generally not performed at that stage. Different diluent extracts of the dried plant parts are generally evaluated for their reported efficacies, though it has been argued that the succulent plant parts contain the highest number and quantity of phytoconstituents than any other stage (Pattanayak 2016, 2020). Such plant medicines traditionally used in their succulent condition can be used in the bio-encapsulated form directly after proper validation of efficacy and calculation of doses (Pattanayak 2021b, 2022).

The earlier reported antidepressant plants have been found to contain phytochemicals with antidepressant activity which include alkaloids, flavonoids, glycosides, phenolics, steroidal saponins, essential oils, resins, tannins, phenols, triterpenoids, sesquiterpenes. The extract of plant parts like roots, leaves, and whole plants when administered in different doses in mice for behavioral experiments, exhibited antidepressant impact (Rahman *et al.* 2017, Martins 2018, de Souza *et al.* 2022). The mechanism of action of most antidepressant plants involves increasing levels of neurotransmitters like Serotonin, Norepinephrine, or Dopamine in the brain, interaction with their receptor systems and corresponding transporters; or through inhibition of Monoamine oxidase enzymes which break down the neurotransmitters by oxidative deamination thereby increasing the concentration of neurotransmitters in the brain (Bondy 2002, Martins 2018, Khushboo *et al.* 2017).

**Table 1. Different antidepressant plants recorded in the literature.**

(Abbreviations used are - SERT: Serotonin Transporter; NE: Norepinephrine; NET: Norepinephrine Transporter; DAT: Dopamine Transporter; MAO: Monoamine Oxidase; MAO-A: Monoamine Oxidase A; MAO-B: Monoamine Oxidase B; GABA: Gamma-Aminobutyric Acid; 5-HT: 5-hydroxytryptamine; CNS: Central Nervous System; cAMP: Cyclic Adenosine Monophosphate; PPAR gamma: Peroxisome proliferator-activated receptor gamma).

Scientific name [Family]	Nutraceutical compound reported	Effects on	Reference
<i>Actaea racemosa</i> L. [Ranunculaceae]	--	Partial agonist of serotonin receptor subtypes	Khushboo <i>et al.</i> 2017, Burdette <i>et al.</i> 2003
<i>Acorus tatarinowii</i> Schott. [Araceae]	--	Antidepressant activity	Li <i>et al.</i> 2001, 2004
<i>Aegle marmelos</i> (L.) Correa [Rutaceae]	--	Increase monoamines level at post-synaptic site; anxiolytic and antidepressant activity	Rahman <i>et al.</i> 2017
<i>Agapanthus campanulatus</i> F.M. Leight [Agapanthaceae]	Flavonoids	Inhibition of SERT, NET and DAT	Martins 2018, Pederson <i>et al.</i> 2008
<i>Akebia quinata</i> (Houtt.) Decne [Lardizabalaceae]	Hederagenin	Inhibition of SERT, NET and DAT	Martins 2018
<i>Alafia multiflora</i> (Stapf) Stapf [Apocynaceae]	Flavonoids, tannins, alkaloids, phenols	Antioxidant, effect on CNS, antidepressant activity	Rahman <i>et al.</i> 2017
<i>Albizia julibrissin</i> Durazz [Fabaceae]	Saponins	Activation of the serotonergic system	Martins 2018, Kim <i>et al.</i> 2007
<i>Allium cepa</i> L. [Amaryllidaceae]	Quercetin glycosides (Flavonoids)	Inhibitor of Monoamine oxidase (MAO)	Martins 2018, Sakakibara <i>et al.</i> 2008
<i>Allium macrostemon</i> Bunge [Amaryllidaceae]	--	Behavioral despair	Khushboo <i>et al.</i> 2017
<i>Allium sativum</i> L. [Amaryllidaceae]	--	Behavioral despair	Khushboo <i>et al.</i> 2017
<i>Aloysia polystachya</i> (Griseb.) Moldenke [Verbenaceae]	--	Antidepressant activity	Khushboo <i>et al.</i> 2017, Martins 2018
<i>Andrographis paniculata</i> (Burm.f.) Nees [Acanthaceae]	--	Antidepressant activity	Rahman <i>et al.</i> 2017
<i>Anemarrhena asphodeloides</i> Bunge [Asparagaceae]	Sarsasapogenin	Inhibition of MAO-A and MAO-B activities	Martins 2018 Ren <i>et al.</i> 2006
<i>Aniba riparia</i> Nees (Mez) [Lauraceae]	Riparin II, Riparin III	Interaction with serotonergic, noradrenergic and dopaminergic receptor systems	Teixeira <i>et al.</i> 2013, Sousa <i>et al.</i> 2004
<i>Apocynum venetum</i> Linn. [Apocynaceae]	Hyperoside, Isoquercitrin	Increase in NE and Dopamine levels in the hippocampus	Martins 2018, Khushboo <i>et al.</i> 2017
<i>Areca catechu</i> L. [Arecaceae]	Saponins	Increase in 5-HT and NE levels in the hippocampus	Dar <i>et al.</i> 1997, 2000
<i>Artemisia absinthium</i> L. [Asteraceae]	--	Antidepressant activity	Ahangar <i>et al.</i> 2011, Martins 2018
<i>Asparagus racemosus</i> Willd. [Asparagaceae]	Steroidal saponins, essential oils, resins, tannins, glycosides	Interaction with Serotonergic, Noradrenergic, Dopaminergic, GAB Aergic receptor systems	Dhingra and kumar 2007, Singh <i>et al.</i> 2009
<i>Bacopa monnieri</i> (L.) Pennell [Plantaginaceae]	--	Inhibition of MAO-A and MAO-B activities	Girish <i>et al.</i> 2016, Martins 2018

Antidepressant plants and their use by the Chiru Tribes of Manipur, India

Scientific name [Family]	Nutraceutical compound reported	Effects on	Reference
<i>Basella alba</i> L. [Basellaceae]	Flavonoids, saponins, phenolic and tannins	Antidepressant activity	Rahman <i>et al.</i> 2017
<i>Berberis aristata</i> DC. [Berberidaceae]	Berberine (Alkaloid)	CNS, Inhibit Monoamine oxidase	Khushboo <i>et al.</i> 2017
<i>Benincasa hispida</i> (Thunb.) Cogn. [Cucurbitaceae]	--	Interaction with Serotonergic, Noradrenergic, Dopaminergic and GAB Aergic receptor systems	Dhingra <i>et al.</i> 2012a, Martins 2018
<i>Boophone disticha</i> (L.f.) [Amaryllidaceae]	Buphanidrine, Buphanamine (Alkaloids)	Inhibition of SERT, NET and DAT	Martins 2018, Pedersen <i>et al.</i> 2008
<i>Bupleurum falcatum</i> L. [Apiaceae]	--	Psycho stimulant effect	Khushboo <i>et al.</i> 2017
<i>Camellia sinensis</i> (L.) Kuntze [Theaceae]	Theanine	Antidepressant activity	Marmat <i>et al.</i> 2013, Singal <i>et al.</i> 2006
<i>Canavalia brasiliensis</i> Benth. [Leguminosae]	Lectins	Interaction with serotonergic, noradrenergic and dopaminergic receptor systems	Martins 2018, Barauna <i>et al.</i> 2006
<i>Cannabis sativa</i> L. [Cannabaceae]	Cannabinoids	Enhanced efflux of Noradrenaline, 5- HT and Dopamine levels	de Souza <i>et al.</i> 2022
<i>Carthamus tinctorius</i> L. [Asteraceae]	--	Antidepressant activity	Martins 2018, Maleki <i>et al.</i> 2013
<i>Casimiroa edulis</i> La Llave [Rutaceae]	Hydroalcoholic extract	Antidepressant activity	Martins 2018, Mora <i>et al.</i> 2005
<i>Cayratia japonica</i> (Thunb.) Gagnep. [Vitaceae]	Flavonoids	MAO inhibition	Martins 2018, Han <i>et al.</i> 2007
# <i>Centella asiatica</i> (L.) [Apiaceae]	Triterpines, saponins, flavonoids, phenolic acids	Increase levels of Monoamine neurotransmitters.	Pardhe <i>et al.</i> 2020, Rahman <i>et al.</i> 2017
<i>Cissampelos sympodialis</i> Eichl. [Menispermaceae]	Total tertiary alkaloids	Antidepressant activity	Martins 2018, Netto <i>et al.</i> 2008
<i>Citrus maxima</i> (Burm.) Merr. [Rutaceae]	--	Antidepressant activity	Rahman <i>et al.</i> 2017
<i>Citrus paradisi</i> Macfad. [Rutaceae]	--	Antidepressant activity	Martins 2018, Gupta <i>et al.</i> 2010.
<i>Clitoria ternatea</i> Linn. [Fabaceae]	--	Cognitive behaviour, anxiety, depression, stress	Martins 2018, Parvathi <i>et al.</i> 2013
<i>Coleus forskohlii</i> Briq. [Lamiaceae]	Forskolin	Increasing brain cAMP availability	Maeda <i>et al.</i> 1997, Martins 2018
<i>Convolvulus pluricaulis</i> Choisy. [Convulvulaceae]	--	Restores brain monoamines	Dhingra <i>et al.</i> 2007, Martins 2018
<i>Crocus sativus</i> L. [Iridaceae]	Ethanollic extract, Kaempferol, Safranal,	Induces dopamine and glutamate release in the brain	Hosseinzadeh <i>et al.</i> 2007, Martins 2018
<i>Cucurbita pepo</i> L. [Cucurbitaceae]	Flavonoids, glycosides	Antioxidant and Antidepressant activity (protects brain feature from CNS disturbance)	Pardhe <i>et al.</i> 2020
# <i>Curcuma longa</i> L. [Zingiberaceae]	Volatile oils-turmerone, curcuminoids, curcumin, sesquiterpenes	Curcumin inhibits MAO enzymes; expands serotonin, noradrenalin and dopamine level in brain	Pardhe <i>et al.</i> 2020, Martins 2018

Scientific name [Family]	Nutraceutical compound reported	Effects on	Reference
<i>Dracocephalum moldavica</i> L. [Lamiaceae]	Triterpenoids, steroids, flavonoids, alkaloids, phenols, coumarins	Antidepressant activity	Pardhe <i>et al.</i> 2020
<i>Echium amoenum</i> Fisch. & C.A. Mey [Boraginaceae]	--	Antidepressant activity	Martins 2018, Sayyah <i>et al.</i> 2009
<i>Eleutherococcus senticosus</i> (Rupr. & Maxim) Maxim. [Araliaceae]	--	Antidepressant activity	Martins 2018, Kimura <i>et al.</i> 2004
<i>Epimedium brevicornum</i> Maxim. [Berberidaceae]	Icariin (Flavonoid)	Antidepressant activity	Martins, 2018, Pan <i>et al.</i> 2007
<i>Galphimia glauca</i> Cav. [Malpighiaceae]	--	Antidepressant activity	Martins 2018 Ruiz <i>et al.</i> 2006
<i>Gastrodia elata</i> Blume [Orchidaceae]	--	Regulates both serotonergic and dopaminergic systems	Martins 2018, Hong <i>et al.</i> 2013
<i>Gentiana kochiana</i> E.P. Perrier & Sonjeon [Gentianaceae]	Xanthones (Gentiacaulein)	Inhibition of MAO-A activity	Martins 2018, Tomic <i>et al.</i> 2005
<i>Ginkgo biloba</i> L. [Ginkgoaceae]	--	Antistress, antidepressant activity	Khushboo <i>et al.</i> 2017
<i>Glycyrrhiza glabra</i> L. [Fabaceae]	Triterpenoidal saponins, flavonoids, alkaloids, phenolic acids	Inhibition of MAO-A activity, increase in brain Norepinephrine and Dopamine levels	Pardhe <i>et al.</i> 2020, Martins 2018
<i>Glycyrrhiza uralensis</i> Fisch. Ex DC. [Fabaceae]	Liquiritin, Isoliquiritin (Flavonoids)	Increase 5-HT and NE in the hippocampus, hypothalamus and cortex	Fan <i>et al.</i> 2012, Zhaoa <i>et al.</i> 2008
<i>Gossypium herbaceum</i> L. [Malvaceae]	--	Activates adenylyl cyclase-cAMP pathway in signal transduction system	Martins 2018, Khaleequr <i>et al.</i> 2012
<i>Hippeastrum vittatum</i> (L' Her) [Amaryllidaceae]	Montanine (Alkaloid)	Neurological disorders and Neurodegenerative disease	Da Silva <i>et al.</i> 2006, Martins 2018
<i>Humulus lupulus</i> L. [Cannabaceae]	Alpha-acids	Antidepressant activity	Martins 2018, Zanolli <i>et al.</i> 2005
<i>Hypericum canariense</i> L. [Hypericaceae]	Methanolic Extract	Neuro pharmacological effect, Anti-cholinergic and sedative properties	Mateo <i>et al.</i> 2005, Khushboo <i>et al.</i> 2017
<i>Hypericum glandulosum</i> Aiton [Hypericaceae]	Methanolic Extract	Neuro pharmacological effect, Anti-cholinergic and sedative properties	Mateo <i>et al.</i> 2005, Khushboo <i>et al.</i> 2017
<i>Hypericum caprifoliatum</i> Cham. & Schltldl. [Hypericaceae]	Phloroglucinol derivatives	Monoamine uptake inhibition	Viana <i>et al.</i> 2005, Martins 2018
<i>Hypericum perforatum</i> L. and <i>Hypericum maculatum</i> Crantz. [Hypericaceae]	Hyperforin	Antidepressant activity	Gambarana <i>et al.</i> 2001, Rahman <i>et al.</i> 2017
<i>Hypericum reflexum</i> L.f. [Hypericaceae]	--	Effect on CNS, Antidepressant activity	Mateo <i>et al.</i> 2001
<i>Ilex pubescens</i> Hook. & Arn. [Aquifoliaceae]	--	Antidepressant	Xu <i>et al.</i> 2004
<i>Inula japonica</i> Thunb. [Asteraceae]	Japonicins (Flavonol)	Antidepressant activity	Martins 2018, Yu <i>et al.</i> 2006

Antidepressant plants and their use by the Chiru Tribes of Manipur, India

Scientific name [Family]	Nutraceutical compound reported	Effects on	Reference
<i>Kaempferia parviflora</i> Wall. ex Baker [Zingiberaceae]	--	Psychiatric illness	Martins 2018, Hawiset <i>et al.</i> 2011
<i>Kielmeyera coriacea</i> Mart. Et Zucc. [Calophyllaceae]	Xanthones	Monoamine uptake inhibition	Martins 2006, Goulart <i>et al.</i> 2007
<i>Lafoensia pacari</i> A. St. Hil. [Lythraceae]	--	Effect on CNS, Antidepressant activity	Martins 2018, Galdino <i>et al.</i> 2009
<i>Lavandula angustifolia</i> Mill. [Lamiaceae]	Linalool, Linalyl acetate	Antidepressant activity	Kageyama <i>et al.</i> 2012, Hritcu <i>et al.</i> 2012
<i>Lepidium meyenii</i> Walp. [Brassicaceae]	--	Antidepressant activity	Martins 2018, Rubio <i>et al.</i> 2006
# <i>Magnolia champaca</i> (L.) Baill. ex Pierre [Magnoliaceae]	Bioactive compounds	Antidepressant, anxiolytic properties	Pushpa <i>et al.</i> 2022
<i>Magnolia officinalis</i> Rehder & Wilson [Magnoliaceae]	Magnolol, Honokiol	GABA modulators, agonist to PPAR gamma	Nakazawa <i>et al.</i> 2003a, Xu <i>et al.</i> 2008
<i>Malus domestica</i> Borkh. [Rosaceae]	Flavonoids, phenolic compounds	Antioxidant rich, involves in growth and production of neurotransmitter Acetylcholine	Pardhe <i>et al.</i> 2020
# <i>Marsilea minuta</i> Linn. [Marsileaceae]	Ethanollic extract, Marsiline (sedative)	Antidepressant activity	Bhattamisra <i>et al.</i> 2008
<i>Melissa officinalis</i> L. [Lamiaceae]	--	Similar activity like antidepressant drug imipramine, strengthens memory, relieves stress-induced headache	Taiwo <i>et al.</i> 2012
<i>Mimosa pudica</i> L. [Fabaceae]	--	Act as Tricyclic Antidepressants	Molina <i>et al.</i> 1999
<i>Mitragyna speciosa</i> (Korth.) Havil. [Rubiaceae]	Mitragynine	Antidepressant activity	Martins 2018, Idayu <i>et al.</i> 2011
<i>Momordica charantia</i> L. [Cucurbitaceae]	Alkaloids, flavonoids, glycosides, steroids, phenols	Dependent on serotonergic, noradrenergic, dopaminergic receptor systems	Ishola <i>et al.</i> 2013
<i>Mondia whitei</i> (Hook.f.) Skeels [Apocynaceae]	--	Affinity for SERT	Pedersen <i>et al.</i> 2008
<i>Morinda officinalis</i> L. [Rubiaceae]	Succinic acid, Nystose, 1F- fructo-furanosylnystose, Inulin-type hexasaccharide, heptasaccharide	Binds selectively to Serotonin receptor	Zhang <i>et al.</i> 2002
<i>Morus alba</i> L. [Moraceae]	--	Antidepressant activity	Sattayasai <i>et al.</i> 2008, Anand <i>et al.</i> 2011
<i>Myristica fragrans</i> Houtt. [Myristicaceae]	--	Antidepressant activity	Dhingra <i>et al.</i> 2006, Moinuddin <i>et al.</i> 2012
<i>Nardostachys jatamansi</i> (D. Don) DC. [Caprifoliaceae]	--	Inhibition of MAO activity, Interaction with GABA receptors	Deepa <i>et al.</i> 2013, Dhingra and Goyal 2008a
<i>Nelumbo nucifera</i> Gaertn. [Nelumbonaceae]	Neferine (Alkaloid)	Serotonin receptor agonist	Martins 2018, Dhanarasu and Al Hazimi 2013

Scientific name [Family]	Nutraceutical compound reported	Effects on	Reference
# <i>Ocimum tenuiflorum</i> L. [Lamiaceae]	Phenolic compounds	Interaction with dopaminergic system	Chatterjee <i>et al.</i> 2011, Pemminati <i>et al.</i> 2010
<i>Paeonia lactiflora</i> Pall. [Paeoniaceae]	Paeoniflorin, Albiflorin (Glycosides)	Effect on Central Monoaminergic Neurotransmitter system, Inhibition of MAO activity	Mao <i>et al.</i> 2008 Rahman <i>et al.</i> 2017
<i>Passiflora foetida</i> L. [Passifloraceae]	Flavonoids, hydrocyanic acid and Harman alkaloids	Effect on CNS disorders, Antidepressant activity	
<i>Paullinia cupana</i> Kunth [Sapindaceae]	Methylxanthines and possibly others	Antidepressant activity	Campos <i>et al.</i> 2005, Martins 2018
<i>Perilla frutescens</i> (L.) Britt [Lamiaceae]	Apigenin, 2,4,5-trimethoxycinnamic acid, Rosemarinic acid	Via dopaminergic mechanism, Cell proliferation in dentate gyrus	Ito <i>et al.</i> 2008, Nakazawa <i>et al.</i> 2003b
# <i>Phyllanthus emblica</i> L. [Phyllanthaceae]	Tannins, gallic acid, alkaloids, phenols	Inhibition of MAO-A activity	Dhingra <i>et al.</i> 2012b, Pemminati <i>et al.</i> 2010
<i>Piper laetispicum</i> C. DC [Piperaceae]	Laetispicine, Leatispiamide A (Alkaloids)	Effect on pain and depression, Antinociceptive properties	Xie <i>et al.</i> 2011, Pan <i>et al.</i> 2005
<i>Piper longum</i> L. [Piperaceae]	Ethanollic extract, Piperine, Methylpiperate	Inhibition of MAO-A and MAO-B activities	Martins 2018, Lee <i>et al.</i> 2005
<i>Piper methysiticum</i> Forst. [Piperaceae]	Pyrones, Desmethoxy yangonin	Inhibition of MAO-B activity, via Dopaminergic mechanism	Martins 2018, Baum <i>et al.</i> 1998
<i>Piper tuberculatum</i> Jacq. [Piperaceae]	Piplartine (Alkaloid)	Anxiolytic and antidepressant activities	Felipe <i>et al.</i> 2007
<i>Plantago asiatica</i> L. [Plantaginaceae]	--	Antidepressant	Xu <i>et al.</i> 2004
<i>Polygala sabulosa</i> A.W. Benn. [Polygalaceae]	Scopoletin, A coumarin	Serotonergic, Dopaminergic and Noradrenergic activities	Khushboo <i>et al.</i> 2017
<i>Polygala tenuifolia</i> Willd. [Polygalaceae]	Polygalatenosides A and B	Norepinephrine reuptake inhibitors, Inhibition of MAO-A and MAO-B activities	Cheng <i>et al.</i> 2006, Hu <i>et al.</i> 2011
<i>Pontederia crassipes</i> Mart. [Pontederiaceae]	Flavonoids, alkaloids, tannins, saponins	Nervine tonic, stimulant, antioxidant, antidepressant	Rahman <i>et al.</i> 2017
<i>Protium heptaphyllum</i> (Aubl.) Marchand [Burseraceae]	Alpha- and Beta-amyrin (Triterpenes)	Antidepressant activity	Aragao <i>et al.</i> 2006, Martins 2018
<i>Psoralea corylifolia</i> L. [Fabaceae]	Psoralen, Psoralidin (Furocoumarins)	Inhibition of MAO-A and MAO-B activities	Martins 2018, Xu <i>et al.</i> 2008
<i>Ptychopetalum olacoides</i> Benth. [Olacaceae]	--	Via D1 dopamine and $\beta$ -noradrenergic receptor systems	Siqueira <i>et al.</i> 2004
<i>Pueraria montana</i> (Lour.) Merr. var. <i>lobata</i> (Willd.) [Fabaceae]	--	Antidepressant activity	Yana <i>et al.</i> 2004, Martins 2018
<i>Rhazya stricta</i> Decne. [Apocynaceae]	--	Effect on Monoamine Oxidase inhibition	Ali <i>et al.</i> 1998, Khushboo <i>et al.</i> 2017
<i>Rhodiola rosea</i> L. [Crassulaceae]	Rosaridin	Via Serotonergic system, Inhibition of MAO-A and MAO-B activities	Mannucci <i>et al.</i> 2012



Antidepressant plants and their use by the Chiru Tribes of Manipur, India

Scientific name [Family]	Nutraceutical compound reported	Effects on	Reference
<i>Rosa damascena</i> Herrm. [Rosaceae]	--	Antidepressant activity	Martins 2018, Rahman <i>et al.</i> 2017
<i>Rosmarinus officinalis</i> L. [Lamiaceae]	Ursolic acid (Triterpenoid), Carnosol, Betulinic acid	Interacts with Monoaminergic system; improves dopaminergic, serotonergic, noradrenergic and cholinergic features.	Machado <i>et al.</i> 2013, Mukhtar <i>et al.</i> 2013
# <i>Rotheca serrata</i> (L.) Steane & Mabb. [Verbenaceae]	Flavonoids	Increases levels of norepinephrine and 5-hydroxytryptamine.	Vazhayil <i>et al.</i> 2017
<i>Roystonea regia</i> (Kunth) O.F.Cook [Arecaceae]	Fatty acids	Antidepressant activity	Carbajal <i>et al.</i> 2009, Martins 2018
<i>Salvia divinorum</i> Epling & Jativa [Lamiaceae]	Salvinorin A	Antidepressant activity	Braida <i>et al.</i> 2009, Martins 2018
<i>Salvia elegans</i> Vahl. [Lamiaceae]	--	Antidepressant activity	Khushboo <i>et al.</i> 2017
<i>Schinus molle</i> L. [Anacardiaceae]	Rutin (Flavonoid)	Dependent on Serotonergic, Noradrenergic and Dopaminergic receptors.	Machado <i>et al.</i> 2007, Khushboo <i>et al.</i> 2017
<i>Schizandra chinensis</i> (Turcz.) Baill. [Schisandraceae]	Flavonoids	Interacts with noradrenergic receptors	Wang <i>et al.</i> 2022
<i>Scrophularia ningpoensis</i> Hemsl. [Scrophulariaceae]	--	Antidepressant activity	Xu <i>et al.</i> 2004
<i>Securidaca longepedunculata</i> Fresen [Polygalaceae]	--	Antidepressant activity	Martins 2018, Adebisi <i>et al.</i> 2006
<i>Selaginella bryopteris</i> L. [Sellaginellaceae]	--	Antidepressant activity	Rahman <i>et al.</i> 2017
<i>Senna occidentalis</i> (L.) Link [Fabaceae]	--	Antianxiety and antidepressant activity	Rahman <i>et al.</i> 2017
<i>Senna spectabilis</i> (D.C.) H.S. Irwin & Barneby [Fabaceae]	Iso-6-spectraline	Antidepressant activity	Silva <i>et al.</i> 2011, Martins 2018
<i>Siphocampylus verticillatus</i> (Cham.) G. Don [Campanulaceae]	--	Interaction with Adrenergic, Dopaminergic, Glutamatergic and Serotonergic system	Rodrigues <i>et al.</i> 2002, Khushboo <i>et al.</i> 2017
<i>Sonchus oleraceus</i> L. [Asteraceae]	--	Antidepressant activity	Vilela <i>et al.</i> 2010, Martins 2018
<i>Sphaeranthus indicus</i> Gaertn. [Asteraceae]	--	Anxiety, depression and convulsions	Galani and Patel 2010
<i>Tabebuia avellanedae</i> Lorentz ex Griseb. [Bignoniaceae]	--	Antidepressant activity	Freitas <i>et al.</i> 2013
# <i>Tagetes lucida</i> Cav. [Asteraceae]	--	Inhibits synaptosomal uptake of Serotonin, Noradrenaline and Dopamine	Gabriela <i>et al.</i> 2012
<i>Tecoma stans</i> (L.) Juss.ex Kunth [Bignoniaceae]	Flavonoids	Antidepressant activity	Rahman <i>et al.</i> 2017
<i>Thymus fallax</i> Fisch. & C.A. Mey. [Lamiaceae]	--	Antidepressant activity	Martins 2018, Semnani <i>et al.</i> 2007
<i>Thymus kotschyanus</i> Boiss. & Hohen. [Lamiaceae]	--	Antidepressant activity	Martins 2018, Semnani <i>et al.</i> 2007

Scientific name [Family]	Nutraceutical compound reported	Effects on	Reference
<i>Thymus pubescens</i> Boiss. & Kotschy ex Celak. [Lamiaceae]	--	Antidepressant activity	Martins 2018, Semnani <i>et al.</i> 2007
<i>Tinospora cordifolia</i> (Willd.) Hook.f. & Thomson [Menispermaceae]	--	Inhibits Mao-A and Mao-B activities, Interaction with serotonergic, noradrenergic and dopaminergic receptor systems	Dhingra and Goyel 2008b, Martins 2018
<i>Trichilia catigua</i> A. Juss. [Meliaceae]	--	Inhibition of dopamine and serotonin uptake	Martins 2018, Campos <i>et al.</i> 2005
<i>Trigonella foneum-graceum</i> L. [Fabaceae]	4-hydroxyisoleucine	Via the serotonergic system	Martins 2018, Gaur <i>et al.</i> 2012
<i>Urtica dioica</i> L. [Urticaceae]	Roots and leaves extract	Antidepressant activity like classical antidepressants fluoxetine and haloperidol	Rahman <i>et al.</i> 2017
<i>Valeriana fauriei</i> Briq. [Caprifoliaceae]	Sesquiterpenes	Antidepressant activity	Liu <i>et al.</i> 2012, Oshima <i>et al.</i> 1995
<i>Valeriana officinalis</i> L. [Caprifoliaceae]	--	Mild sleep disorders and Nervous tension, Antidepressant activity	Hattesoehl <i>et al.</i> 2008; Sakamoto <i>et al.</i> 1992, Martins 2018
<i>Valeriana wallichii</i> DC. [Valerianaceae]	--	Nitric oxide inhibition	Subhan <i>et al.</i> 2010, Sah <i>et al.</i> 2011
<i>Vitis vinifera</i> L. [Vitaceae]	--	Antidepressant activity	Martins 2018, Sreemantula <i>et al.</i> 2005
<i>Withania somnifera</i> (L.) Dunal [Solanaceae]	Bioactive glycowithanolides	Anxiolytic and antidepressant action	Bhattacharya <i>et al.</i> 2020
<i>Xysmalobium undulatum</i> (L.) W.T. Aiton [Apocynaceae]	--	Affinity for SERT	Martins 2018, Pedersen <i>et al.</i> 2008
<i>Zingiber officinale</i> Roscoe [Zingiberaceae]	Gingerol, Shogool	Binding to Serotonin receptor	Martins 2018

# Plants are identified in the present study for alike activities.

\$ The sign (--) in the table indicates no known phytochemicals responsible for antidepressant activity.

## CONCLUSION

People living in remote and hilly places like the Chiru tribe depend on their surrounding forests and vegetation for their livelihood- food, fodder, and medicines. The majority of the population depends on traditional knowledge of medicines as modern healthcare facilities are not easily accessible in these remote areas. Besides, ethnomedicinal plants are easily available, reliable, cost-effective, and safe with no undesired side-effects. Also, there is an increasing trend in the use of herbal remedies and plant-based products to treat depression and related mental disorders worldwide because of their great efficacy. People also have stronger trust in their age-old practice which has been passed over generations.

Nutraceuticals in plant-based food have neuroprotective functions and can be prototypes for antidepressant drugs in the future. In this direction, the present finding will form baseline data for developing a standard protocol for the production of herbal antidepressant drugs. The mechanism of understanding many antidepressant plants still needs to be explored to fill the gaps in depression research and to develop better treatment measures. It is therefore crucial to document the ethnomedicinal knowledge as changes in lifestyle and occupation raise a serious concern for the indigenous knowledge, if not documented properly. Also, increasing encroachment of the forests and environmental degradation either due to natural or man-made calamities pose a serious threat to the

## Antidepressant plants and their use by the Chiru Tribes of Manipur, India

medicinal plant diversity. Amidst the rising trend of excessive harvesting of medicinal plants to develop herbal remedies and drugs, it is vital to determine the availability status of these plants and initiate appropriate conservation efforts. It is the need of the hour to conserve selected medicinal plants sustainably for further productive research work.

### ACKNOWLEDGEMENT

The authors acknowledged all the key informants of the *Chiru* tribe who inhabited villages of Manipur for sharing their ethnomedicinal knowledge and ideas. Also, the authors are deeply thankful to all the headmen and the village members for their cooperation and hospitality extended throughout the study. The authors are also thankful to the editorial reviewers for their constructive suggestions.

### REFERENCES

- Adebisi RA, Elsa AT, Agaie BM, Etuk EU (2006) Antinociceptive and antidepressant like effects of *Securidaca longepedunculata* root extract in mice. *J Ethnopharmacol* 107: 234-239.
- Ahangar N, Mirfetros S, Ebrahimzadeh MA (2011) Antidepressant activity of polyphenol fraction of *Artemisia absinthium* L. *Pharmacologyonline* 1: 825-832.
- Ahmad A, Khan A, Safdar M (2011) Curcumin a natural product of pharmaceutical importance. *Int J Pharma Clin Exp Bio* 5(1): 44-49.
- Ali BH, Bashir AK, Tanira MOM (1998) The effect of *Rhazya stricta* Decne, a traditional medicinal plant, on the forced swimming test in rats. *Pharmacol Biochem Behav* 59: 547-550.
- Aragao GF, Carneiro LMV, Junior APF, Vieira LC, Bandeira PN *et al.* (2006) A possible mechanism for anxiolytic and anti-depressant effects of alpha- and beta-amyrin from *Protium heptaphyllum* (Aubl.) March. *Pharmacol Biochem Behav* 85: 827-834.
- Barauna SC, Kaster MP, Heckert BT, Nascimento KSD, Rossi FM *et al.* (2006) Antidepressant-like effect of lectin from *Canavalia brasiliensis* (ConBr) administered centrally in mice. *Pharmacol Biochem Behav* 85: 160-169.
- Baum SS, Hill R, Rommelspacher H (1998) Effect of kava extract and individual kavapyrones on neurotransmitter levels in the nucleus accumbens of rats. *Prog Neuropsychopharmacol Biol Psychiatry* 22: 1105-1120.
- Bhattacharya SK, Bhattacharya A, Sairam K, Ghosal S (2020) Anxiolytic-antidepressant activity of *Withania somnifera* glycowithanolides: an experimental study. *Phytomedicine* 7(6): 463-469. DOI: 10.1016/S0944-7113(00)80030-6.
- Bhattamisra SK, Khannab VK, Agrawal AK, Singh PN, Singh SK (2008) Antidepressant activity of standardised extract of *Marsilea minuta* Linn. *J Ethnopharmacol* 117: 51-57.
- Bondy B (2002) Pathophysiology of depression and mechanisms of treatment. *Dialogues Clinical Neurosci* 4(1): 1-19.
- Braida D, Capurro V, Zani A, Rubino T, Vigano D *et al.* (2009) Potential anxiolytic- and antidepressant-like effects of salvinatorin A, the main active ingredient of *Salvia divinorum*, in rodents. *Br J Pharmacol* 157: 844-853.
- Burdette JE, Liu J, Chen SN, Fabricant DS, Piersen CE *et al.* (2003) Black cohosh acts as a mixed competitive ligand and partial agonist of the serotonin receptor. *J Agric Food Chem* 51: 5661-5670.
- Campos AR, Barros AIS, Albuquerque FAA, Leal LKAM, Rao VSN (2005) Acute effects of guarana (*Paullinia cupana* Mart.) on mouse behaviour in forced swimming and open field tests. *Phytother Res* 19: 441-443.
- Campos MM, Fernandes ES, Ferreira J, Santos ARS, Calixto JB (2005) Antidepressant-like effects of *Trichilia catigua* (Catuaba) extract: evidence for dopaminergic-mediated mechanisms. *Psychopharmacology* 182: 45-53.
- Carbajal D, Ravelo Y, Molina V, Mas R, Arruzazabala MD (2009) D-004, a lipid extract from royal palm fruit, exhibits antidepressant effects in the forced swim test and the tail suspension test in mice. *Pharmacol Biochem Behav* 92: 465-468.
- Chatterjee M, Verma P, Maurya R, Palit G (2011) Evaluation of ethanol leaf extract of *Ocimum tenuiflorum* in experimental models of anxiety and depression. *Pharm Biol* 49: 477-483.
- Cheng MC, Li CY, Ko HC, Ko FN, Lin YL, Wu TS (2006) Antidepressant principles of the roots of *Polygala tenuifolia*. *J Nat Prod* 69: 1305-1309.
- Dar A, Khatoon S, Rahman G, A. Rahman AU (1997) Anti-depressant activities of *Areca catechu* fruit extract. *Phytomedicine* 4: 41-45, DOI: 10.1016/S0944-7113(97)80026-8.
- Dar A, Khatoon S (2000) Behavioral and biochemical studies of dichloromethane fraction from the *Areca catechu* nut. *Pharmacol Biochem Behav* 65: 1-6, DOI: 10.1016/S0091-3057(99)00179-3.

DaSilva AFS, DeAndrade JP, Bevilaqua LRM, DeSouza MM, Izquierdo I *et al.* (2006) Anxiolytic, antidepressant and anticonvulsant-like effects of the alkaloid montanine isolated from *Hippeastrum vittatum*. *Pharmacol Biochem Behav* 85: 148-154.

de Souza MR, Henriques AT, Limberger RP (2022) Medical cannabis regulation: An overview of models around the world with emphasis in the Brazilian scenario. *J Cannabis Res* 4(33): 1-15.

Deepa B, Suchetha K, Rao S (2013) Antidepressant activity of *Nardostachys jatamansi* in electron beam irradiated mice. *Int J Res Ayur Pharm* 4: 101-103.

Dhanarasu S, Al-Hazimi A (2013) Phytochemistry, pharmacological and therapeutic applications of *Nelumbo nucifera*. *Asian J Phytomed Clin Res* 1: 123-136.

Dhingra D, Sharma A (2006) Antidepressant-like activity of n-hexane extract of nutmeg (*Myristica fragrans*) seeds in mice. *J. Med Food* 9: 84-89.

Dhingra D, Joshi P (2012a) Antidepressant-like activity of *Benincasa hispida* fruits in mice: possible involvement of monoaminergic and GABAergic systems. *J Pharmacol Pharmacother* 3: 60-62.

Dhingra D, Joshi P, Gupta A, Chhillar R (2012b) Possible involvement of mono- aminergic neurotransmission in antidepressant-like activity of *Embllica officinalis* fruits in mice. *CNS Neurosci Ther* 18(5): 419-425. DOI: 10.1111/j.1755-5949.2011.00256.x.

Dhingra P, Goyal PK (2008a) Inhibition of MAO and GABA: probable mechanisms for antidepressant-like activity of *Nardostachys jatamansi* DC. in mice. *Indian J Exp Biol* 46: 212-218.

Dhingra P, Goyal PK (2008b) Evidences for the involvement of monoaminergic and GABAergic systems in antidepressant-like activity of *Tinospora cordifolia* in mice. *Indian J Pharm Sci* 70: 761-767, DOI: 10.4103/0250-474X.49118.

Dhingra D, Valecha R (2007) Screening for antidepressant-like activity of *Convolvulus pluricaulis* Choisy in mice. *Pharmacologyonline* 1: 262-278.

Dhingra D, Kumar V (2007) Pharmacological evaluation for antidepressant-like activity of *Asparagus racemosus* Willd. in mice. *Pharmacologyonline* 3: 133-152.

Fan ZZ, Zhao WH, Guo J, Cheng RF, Zhao JY *et al.* (2012) Antidepressant activities of flavonoids from *Glycyrrhiza uralensis* and its neurogenesis protective effect in rats. *Yao Xue Xue Bao* 47: 1612-1617.

Felipe FCB, Filho JLTS, Souza LED, Silveira JA, Uchoa DED *et al.* (2007) Piplartine, an amide alkaloid from *Piper tuberculatum*, presents anxiolytic and antidepressant effects in mice. *Phytomedicine* 14: 605-612.

Freitas AE, Machado DG, Budni J, Neis VB, Balen GO, Lopes MW *et al.* (2013) Antidepressant-like action of the bark ethanolic extract from *Tabebuia avellanedae* in the olfactory bulbectomized mice. *J Ethnopharmacol* 145: 737-745.

Gabriela GC, Javier AAF, Elisa VA, Gonzalo VP, Herlinda BJ (2012) Antidepressant-like effect of *Tagetes lucida* Cav. extract in rats: involvement of the serotonergic system. *Am J Chin Med* 40: 753-768.

Galdino PM, Nascimento MVM, Sampaio BL, Ferreira RN, Paula JR *et al.* (2009) Antidepressant-like effect of *Lafoensia pacari* A. St.-Hil. ethanolic extract and fractions in mice. *J Ethnopharmacol* 124: 581-585.

Gambarana C, Tolu PL, Masi F, Rinaldi M, Giachetti D *et al.* (2001) A study of the antidepressant activity of *Hypericum perforatum* on animal models. *Pharmacopsychiatry* 34: 42-44.

Galani VJ, Patel BG (2010) Effect of hydroalcoholic extract of *Sphaeranthus indicus* against experimentally induced anxiety, depression and convulsions in rodents. *Int J Ayurveda Res* 1(2): 87-92. DOI: 10.4103/0974-7788.64412.

Gaur V, Bodhankar SL, Mohan V, Thakurdesai P (2012) Antidepressant-like effect of 4-hydroxyisoleucine from *Trigonella foenum graecum* L. seeds in mice. *Biomedic Aging Pathol* 2: 121-125.

Girish C, Oommen S, Vishnu R (2016) Evidence for the involvement of the mono- aminergic system in the antidepressant-like activity of methanolic extract of *Bacopa monnieri* in albino mice. *Int J Basic Clin Pharmacol* 3: 914-922.

Goulart YCF, Martins JVC, Santos AR, Moreira LY, Calixto JB *et al.* (2007) Involvement of serotonin in the antidepressant-like effect of extract from *Kielmeyera coriacea* stems. *Pharmaceutic Biol* 45(3): 169-175, DOI: 10.1080/13880200701212981.

Gupta V, Bansal P, Kumar P, Shri R (2010) Anxiolytic and antidepressant activities of different extracts from *Citrus paradisi* var. Duncan. *Asian J Pharm Clin Res* 3: 98-100.

Han XH, Hong SS, Hwang JS, Lee MK., Hwang BY *et al.* (2007) Monoamine oxidase inhibitory components from *Cayratia japonica*. *Arch Pharm Res* 30: 13-17.

Hattesoehl M, Feistel B, Sievers H, Lehnfeld R, Hegger M, Winterhoff H (2008) Extracts of *Valeriana officinalis* L.

## Antidepressant plants and their use by the Chiru Tribes of Manipur, India

- s.l. show anxiolytic and antidepressant effects but neither sedative nor myorelaxant properties. *Phytomedicine* 15: 2-15.
- Hawiset T, Muchimapura S, Wattanathorn J, Sripanidkulchai B (2011) Screening neuropharmacological activities of *Kaempferia parviflora* (Krachai Dam) in healthy adult male rats. *Am J Appl Sci* 8: 695-702.
- Hong SS, Cho SH (2013) Antidepressant-like effects of the *Gastrodia elata* BL extract in mice. *J Orient Neuropsychiatry* 24: 281-292.
- Hosseinzadeh H, Motamedshariaty V, Hadizadeh F (2007) Antidepressant effect of kaempferol, a constituent of saffron (*Crocus sativus*) petal, in mice and rats. *Pharmacologyonline* 2: 367-370.
- Hritcu L, Cioanca O, Hancianu M (2012) Effects of lavender oil inhalation on improving scopolamine-induced spatial memory impairment in laboratory rats. *Phytomedicine* 19: 529-534.
- Hu Y, Liu M, Liu P, Guo DH, Wei RB *et al.* (2011) Possible mechanism of the antidepressant effect of 3,6'-disinapoyl sucrose from *Polygala tenuifolia* Willd. *J Pharm Pharmacol* 63: 869-874.
- Idayu NF, Hidayat MT, Moklas MAM, Sharida F, Raudzah ARN *et al.* (2011) Antidepressant-like effect of mitragynine isolated from *Mitragyna speciosa* Korth in mice model of depression. *Phytomedicine* 18: 402-407.
- Ishola IO, Akinyede AA, Sholarin AM (2013) Antidepressant and anxiolytic properties of the methanolic extract of *Momordica charantia* Linn (Cucurbitaceae) and its mechanism of action. *Drug Res* 64: 368-376.
- Ito N, Yabe T, Gamo Y, Nagai T, Oikawa T *et al.* (2008) Rosmarinic acid from *Perillae herba* produces an antidepressant-like effect in mice through cell proliferation in the hippocampus. *Biol. Pharm. Bull.* 31: 1376-1380.
- Kageyama A, Ueno T, Oshio M, Masuda H, Horiuchi H *et al.* (2012) Antidepressant-like effects of an aqueous extract of Lavender (*Lavandula angustifolia* Mill.) in rats. *J Food Sci Technol* 18: 473-479.
- Khaleequr R, Arshiya S, Shafeequr R (2012) *Gossypium herbaceum* Linn: an ethno- pharmacological review. *J Pharm Sci Innov* 1: 1-5.
- Khushboo, Sharma B (2017) Antidepressants: mechanism of action, toxicity and possible amelioration. *J Appl Biotechnol Bioeng* 3(5): 437-448.
- Kim JH, Kim SY, Lee SY, Jang CG (2007) Antidepressant like effects of *Albizia julibrissin* in mice: involvement of the 5-HT1A receptor system. *Pharmacol Biochem Behav* 87: 41-47.
- Kimura Y, Sumiyoshi M (2004) Effects of various *Eleutherococcus senticosus* cortex on swimming time, natural killer activity and corticosterone level in forced swimming stressed mice. *J Ethnopharmacol* 95: 447-453.
- Lee SA, Hong SS, Han XH, Hwang JS, Oh GJ *et al.* (2005) Piperine from the fruits of *Piper longum* with inhibitory effect on monoamine oxidase and antidepressant-like activity. *Chem Pharm Bull* 53: 832-835.
- Li M, Chen H (2001) Antidepressant effect of water decoction of *Rhizoma acori tatarinowii* in the behavioural despair animal models of depression. *Zhong Yao Cai* 24: 40-41.
- Li M, Li J, Ji N, Guo L, Gan H *et al.* (2004) Antidepressant effect of several crude extracts of *Rhizoma acori tarinowii* (CBA:460369). *Acad J Guangdong Coll Pharm* 20: 141-144.
- Liu XG, Gao PY, Wang GS, Song SJ, Li LZ *et al.* (2012) *In vivo* antidepressant activity of sesquiterpenes from the roots of *Valeriana fauriei* Briq. *Fitoterapia* 83: 599-603.
- Machado DG, Cunha MP, Neis VB, Balen GO, Colla A *et al.* (2013) Antidepressant-like effects of fractions, essential oil, carnosol and betulinic acid isolated from *Rosmarinus officinalis* L. *Food Chem* 136: 999-1005.
- Machado DG, Kaster MP, Binfaré RW, Dias M, Santos AR *et al.* (2007) Antidepressant-like effect of the extract from leaves of *Schinus molle* L. in mice: evidence for the involvement of the monoaminergic system. *Prog Neuropsychopharmacol Biol Psychiatry* 31(2): 421-428, DOI: 10.1016/j.pnpbp.2006.11.004.
- Maeda H, Ozawa H, Saito T, Irie T, Takahata N (1997) Potential antidepressant properties of forskolin and a novel water-soluble forskolin (NKH477) in the forced swimming test. *Life Sci* 61: 2435-2442.
- Maleki SA, Asgarpanah J, Bekhradi R, Maleki FA, Ahangari NM *et al.* (2013) Antidepressant like effect of ethanolic extract of *Carthamus tinctorius* in the forced swim test and tail suspension test in male mice. *Chin J Integr Med* 19: 153-159.
- Mannucci C, Navarra M, Calzavara E, Caputi AP, Calapai G (2012) Serotonin involvement in *Rhodiola rosea* attenuation of nicotine withdrawal signs in rats. *Phytomedicine* 19: 1117-1124.
- Mao QQ, Ip SP, Tsai SH, Che CT (2008) Antidepressant-like effect of peony glycosides in mice. *J Ethnopharmacol* 119: 272-275.

- Marmat A, Middha AK (2013) A study on antidepressant activity of leaves extracts of *Camellia sinensis*. Intl J Pharm Innov 3: 99-110.
- Martin GJ (2007) Ethnobotany-A methods Manual, Earthscan, London.
- Martins J, Otobone FJ, Sela VR, Obici S, Trombelli MA *et al.* (2006) Behavioral effects of *Kielmeyera coriacea* extract in rats. Indian J Pharmacol 38: 427-428.
- Martins JSB (2018) Phytochemistry and pharmacology of anti-depressant medicinal plants: A review. Biomed Pharmacother 104: 343-365, DOI: 10.1016/j.biopha.2018.05.044.
- Mateo CCS, Bonkanka CX, Prado B, Rabanal RM (2005) Antidepressant properties of some *Hypericum canariense* L. and *Hypericum glandulosum* Ait. extracts in the forced swimming test in mice. J Ethnopharmacol 97: 541-547.
- Moinuddin G, Devi K, Khajuria DK (2012) Evaluation of the anti-depressant activity of *Myristica fragrans* (Nutmeg) in male rats. Avicenna J Phytomed 2: 72-78.
- Molina M, Contreras CM, P.T. Alcantara PT (1999) *Mimosa pudica* may possess anti-depressant actions in the rat. Phytomedicine 6: 319-323.
- Mora S, Veliz GD, Lungenstrass H, Gonzalez MG, Morales TC *et al.* (2005) Central nervous system activity of the hydroalcoholic extract of *Casimiroa edulis* in rats and mice. J Ethnopharmacol 97: 191-197.
- Mukhtar EJA, Selman SM, Sahib ZH, Naji H (2013) Antidepressant-like effect of *Rosmarinus officinalis* extract in male mice. Med J Babylon 10: 803-808.
- Nakazawa T, Yasuda T, Ohsawa K (2003a) Metabolites of orally administered *Magnolia officinalis* extract in rats and man and its antidepressant-like effects in mice. J Pharm Pharmacol 55: 1583-1591.
- Nakazawa T, Yasuda T, Ueda J, Ohsawa K (2003b) Antidepressant-like effects of apigenin and 2,4,5-trimethoxycinnamic acid from *Perilla frutescens* in the forced swimming test. Biol Pharm Bull 26: 474-480.
- Netto SM, Varela RWB, Fachine MF, Queiroga MNG, Maior FNS *et al.* (2008) Antidepressant effects of total tertiary alkaloid fraction of *Cissampelos sympodialis* Eichler in rodents. Braz J Pharmacol 18: 165-169.
- Oshima Y, Matsuoka S, Ohizumi Y (1995) Antidepressant principles of *Valeriana fauriei* roots. Chem Pharm Bull (Tokyo) 43: 169-170.
- Pan SL, Xie J, Qian FG, Wang J, Shao YC (2005) Antidepressant amides from *Piper laetispicum* C. DC, Yao Xue Xue Bao 40: 355-357.
- Pan Y, Kong LD, Li YC, Xia X, Kung HF *et al.* (2007) Icaritin from *Epimedium brevicornum* attenuates chronic mild stress-induced behavioral and neuroendocrinological alterations in male Wistar rats. Pharmacol Biochem Behav 87: 130-140.
- Pannu A, Sharma PC, Thakur VK, Goyal RK (2021) Emerging role of flavonoids as the treatment of depression. Biomolecules 11: 1-49.
- Pardhe HA, Nagalakshmi NC, Hariprashad MG, Chourasia PK, Nandini S (2020) A review: medicinal plants with antidepressant properties. IP Indian J Neurosci 6(1): 1-5.
- Parvathi M, Ravishankar K (2013) Evaluation of antidepressant, motor coordination and locomotor activities of ethanolic root extract of *Clitoria ternatea*. J Nat Remedies 13: 19-24.
- Patel A, Shah H, Gandhi T (2022) Saponin rich fraction of *Bauhinia variegata* Linn. ameliorates kidney stone formation in Rats. Explor Anim Med Res 12(1): 74-84. DOI: 10.52635/eamr/12.1.74-84.
- Pattanayak S, Mandal TK, Bandyopadhyay SK (2016) Validation and therapeutic use of succulent plant parts - opening of a new horizon of alternative medicine. Explor Anim Med Res 6(1): 8-14.
- Pattanayak S (2020) Succulent biomedicines - an effective way of getting protection against diseases through immunomodulation. Explor Anim Med Res 10(2): 112-123.
- Pattanayak S (2021a) Plants in healthcare: past, present and future. Explor Anim Med Res 11(2): 140-144. DOI: 10.52635/eamr/11.2.140-144.
- Pattanayak S (2021b) Anti-COVID-19 biomedicines - a layout proposal for production, storage and transportation. The Open COVID J 1: 166-88. DOI: 10.2174/2666958702101010166.
- Pattanayak S (2022) Prevention and control of diabetes by intake of succulent biomedicines and following of designed lifestyle: A ready plan for execution. Internati J Scientific Res Updates 3(2): 081-103. <https://doi.org/10.53430/ijrsru.2022.3.2.0047>.
- Paul A, Sujatha K (2022) Concurrent effect of *Linum usitatissimum* and *Emblica officinalis* on lead induced oxidative stress and histomorphological changes in uterus of female Wistar rats. Explor Anim Med Res 12(2): 264-272, DOI: 10.52635/eamr/12.2.264-272.

## Antidepressant plants and their use by the Chiru Tribes of Manipur, India

- Pedersen ME, Szewczyk B, Stachowicz K, Wieronska J, Andersen J *et al.* (2008) Effects of South African traditional medicine in animal models for depression. *J Ethnopharmacol* 19: 542-548.
- Pemminati S, Gopalakrishna HN, Alva A, Pai MRSM, Seema Y *et al.* (2010) Antidepressant activity of ethanolic extract of leaves of *Ocimum sanctum* in mice. *J Pharm Res* 3: 624-626.
- Pradhan DK, Ghosh J, Lepcha N, Nandi A, Banerjee D *et al.* (2021) New ethnomedicinal information from Lepcha community of Dzongu, Sikkim. *Explor Anim Med Res* 11(2): 179- 187, DOI: 10.52635/eamr/11.2.179-187.
- Pushpa VH, Jayanthi MK, Rashmi HR, Kumar V, Shivamurthy N *et al.* (2022) New insights on the phytochemical intervention for the treatment of neuropsychiatric disorders using the leaves of *Michelia champaca*: an *in vivo* and *in silico* approach, *Pharmaceutic Biol* 60(1): 1656-1668, DOI: 10.1080/13880209.2022.2101669.
- Rahman MR, Ali M, Sharif M, Tajmim A (2017) A review study on the traditional plants has potential antidepressant property. *MOJ Cell Sci Rep* 4(5): 1-9.
- Ren LX, Luo YF, Li X, Zuo DY, Wu YL (2006) Antidepressant-like effects of Sarsasapogenin from *Anemarrhena asphodeloides* Bunge (Liliaceae). *Biol Pharm Bull* 29: 2304-2306.
- Rodrigues ALS, Silva GLD, Mateussi AS, Fernandes ES, Miguel OG *et al.* (2002) Involvement of monoaminergic system in the antidepressant-like effect of the hydroalcoholic extract of *Siphocampylus verticillatus*. *Life Sci* 70: 1347-1358.
- Rubio J, Caldas M, Davila S, Gasco M, Gonzales GF (2006) Effect of three different cultivars of *Lepidium meyenii* (Maca) on learning and depression in ovariectomized mice. *BMC Complement Altern Med* 6: 23.
- Ruiz HM, Ferrer JJE, DeLima TC, Montes AD, Garcia PD *et al.* (2006) Anxiolytic and antidepressant-like activity of a standardized extract from *Galphimia glauca*. *Phytomedicine* 13: 23-28.
- Sah SP, Mathela CS, Chopra K (2011) Antidepressant effect of *Valeriana wallichii* patchouli alcohol chemotype in mice: behavioural and biochemical evidence. *J Ethnopharmacol* 135: 197-200.
- Sakakibara H, Yoshino S, Kawai Y, Terao J (2008) Antidepressant-like effect of onion (*Allium cepa* L.) powder in a rat behavioral model of depression. *Biosci Biotechnol Biochem* 72: 94-100.
- Sakamoto T, Mitani Y, Nakajima K (1992) Psychotropic effects of Japanese valerian root extract. *Chem Pharm Bull (Tokyo)* 40: 758-761.
- Sattayasai J, Tiamkao S, Puapairoj P (2008) Biphasic effects of *Morus alba* leaves green tea extract on mice in chronic forced swimming model. *Phytother Res* 22: 487492.
- Sayyah M, Boostani H, Pakseresht S, Malaieri A (2009) Efficacy of aqueous extract of *Echium amoenum* in treatment of obsessive-compulsive disorder. *Prog Neuropsychopharmacol Biol Psychiatry* 33: 1513-1516.
- Semnani KM, Mahmoudi M, Riahi G (2007) Effects of essential oils and extracts from certain *Thymus* species on swimming performance in mice. *Pharm Biol* 45: 464-467.
- Silva FO, Silva MG, Cerqueira GS, Sabino EB, Almeida AAC (2011) Central nervous system effects of Iso-6-spectraline isolated from *Senna spectabilis* var. *Excelsa* (Schrad) in mice. *J Young Pharm* 3: 232-236.
- Singal A, Tirkey N, Muragundla A, Chopra K (2006) Green tea [*Camellia sinensis* (L.) O. Kuntze] extract reverses the despair behavior in reserpinised and diabetic mice. *Indian J Exp Biol* 44: 913-917.
- Singh GK, Garabadu D, Muruganandam AV *et al.* (2009) Antidepressant activity of *Asparagus racemosus* in rodent models. *Pharmacol Biochem Behav* 91(3): 283-290.
- Siqueira IR, Cimarosti H, Fochesatto C, Nunes DS, Salbego C *et al.* (2004) Neuroprotective effects of *Ptychopetalum olacoides* Benth (Olacaceae) on oxygen and glucose deprivation induced damage in rat hippocampal slices. *Life Sci* 75: 1897-1906.
- Sousa FCF, Melo CTV, Monteiro AP, Lima VTM, Gutierrez SJC *et al.* (2004) Antianxiety and antidepressant effects of riparin III from *Aniba riparia* (Nees) Me (Lauraceae) in mice. *Pharmacol Biochem Behav* 78: 27-33.
- Sreemantula S, Nammi S, Kolanukonda R, Koppula S, Boini KM (2005) Adaptogenic and nootropic activities of aqueous extract of *Vitis vinifera* (grape seed): an experimental study in rat model. *BMC Complement Altern Med* 5: 1, <https://doi.org/10.1186/1472-6882-5-1>.
- Subhan F, Karim N, Gilani AH, Sewell RDE (2010) Terpenoid content of *Valeriana wallichii* extracts and antidepressant-like response profiles. *Phytother Res* 24: 686-691.
- Taiwo AE, Leite FB, Lucena GM, Barros M, Silveira D *et al.* (2012) Anxiolytic and antidepressant like effects of *Melissa officinalis* (lemon balm) extract in rats: influence of administration and gender. *Indian J Pharmacol* 44: 189-192.

Teixeira CP, DeMelo CT, DeAraujo FL, DeCarvalho AM, Silva MI *et al.* (2013) Antidepressant-like effect of riparin II from *Aniba riparia* in mice: evidence for the involvement of the monoaminergic system. *Fundam Clin Pharmacol* 27: 129-137.

Tomic M, Tovilovic G, Butorovic B, Krstic D, Jankovic T *et al.* (2005) Neuropharmacological evaluation of diethylether extract and xanthenes of *Gentiana kochiana*. *Pharmacol Biochem Behav* 81: 535-542.

Vazhayil BK, Rajagopal SS, Thangavelu T, Swaminathan G, Rajagounder E (2017) Neuroprotective effect of *Clerodendrum serratum* Linn. leaves extract against acute restraint stress-induced depressive-like behavioral symptoms in adult mice. *Indian J Pharmacol* 49(1): 34-41.

Viana A, DoRego JC, Poser VG, Ferraz A, Heckler AP *et al.* (2005) The antidepressant-like effect of *Hypericum caprifoliatum* Cham and Schlecht (Guttiferae) on forced swimming test results from an inhibition of neuronal monoamine uptake. *Neuropharmacol* 49: 1042-1052.

Vilela FC, Padilha MM, Silva GAD, Soncini R, Paiva AG (2010) Antidepressant-like activity of *Sonchus oleraceus* in mouse models of immobility tests. *J Med Food* 13: 219-222.

Wang J, Zhang G, Yang Y, Zhang X, Shi K *et al.* (2022) *Schisandra chinensis* Lignans exert antidepressant effects by promoting BV2 microglia polarization toward the M2 phenotype through the activation of the cannabinoid receptor type-2-signal transducer and activator of transcription 6 pathway. *J Agric Food Chem* 70(44): 14157-14169, <https://doi.org/10.1021/acs.jafc.2c04565>.

Xie H, Yan MC, Jin D, Liu JJ, Yu M (2011) Studies on antidepressant and antinociceptive effects of ethyl acetate extract from *Piper laetispicum* and structure-activity

relationship of its amide alkaloids. *Fitoterapia* 82: 1086-1092.

Xu C, Luo L, Tan RX (2004) Antidepressant effect of three traditional Chinese medicines in the learned helplessness model. *J Ethnopharmacol* 91: 345-349.

Xu Q, Yi LT, Pan Y, Wang X, Li YC *et al.* (2008) Antidepressant-like effects of the mixture of honokiol and magnolol from the barks of *Magnolia officinalis* in stressed rodents. *Prog Neuropsychopharmacol Biol Psychiatry* 32: 715-725, DOI: 10.1016/j.pnpbp.2007.11.020.

Xu Q, Pan Y, Yi LT, Li YC, Mo SF *et al.* (2008) Antidepressant-like effects of Psoralen isolated from the seeds of *Psoralea corylifolia* in the mouse forced swimming test. *Biol Pharm Bull* 31: 1109-1114.

Yana B, Wang DY, Xing DM, Ding Y, Wang RF *et al.* (2004) The anti-depressant effect of ethanol extract of *Radix puerariae* in mice exposed to cerebral ischemia reperfusion. *Pharmacol Biochem Behav* 78: 319-325.

Yu NJ, Zhao YM, Zhang YZ, Li YF (2006) Japonicins A and B from the flowers of *Inula japonica*. *J Asian Nat Prod Res* 8: 385-390.

Zanoli P, Rivasi M, Zavatti M, Brusiani F, Baraldi M (2005) New insight in the neuropharmacological activity of *Humulus lupulus* L. *J Ethnopharmacol* 102: 102-106.

Zhang ZQ, Yuan L, Yang M, Luo ZP, Zhao YM (2002) The effect of *Morinda officinalis*, how a Chinese traditional medicinal plant, on the DRL 72-s schedule in rats and the forced swimming test in mice. *Pharmacol Biochem Behav* 72: 39-43.

Zhaoa Z, Wang W, Guob H, Zhoua D (2008) Antidepressant-like effect of liquiritin from *Glycyrrhiza uralensis* in chronic variable stress induced depression model rats. *Behav Brain Res* 194: 108-113.

**\*Cite this article as:** Chanu SP, Rajkumari R, Pinokiyo A (2023) Antidepressant plants and their use by the *Chiru* tribes of Manipur, India. *Explor Anim Med Res* 13(Ethnomed. Spl.): 51-66, DOI: 10.52635/eamr/13(S)51-66.