

*Research Article*

## SERO-PREVALENCE OF BOVINE LEPTOSPIROSIS IN SOUTH ANANDAMAN ISLANDS, INDIA

Joyjit Mitra<sup>1</sup>, Sumit Chowdhury<sup>2\*</sup>, Shibabrata Pattanayak<sup>3</sup>

**ABSTRACT:** A total of 108 blood serum samples of bovine were collected from different regions of South Andaman to Regional Disease Diagnostic Laboratory (RDDL, ER, Kolkata) for sero-prevalence study of bovine leptospirosis. All the sera samples were tested by Microscopic Agglutination Test (MAT) against eight serovars of *Leptospira* spp. Out of 108 serum samples tested, 75(69.44%) were serologically positive. The most prevalent serovars were Automnalis (53.70%) followed by Sejroe (28.70%) and Hardjo (22.22%). This study suggests that bovine may have a role in maintaining Automnalis serovar of *Leptospira* as reservoir in Anandaman and Nicobar Islands.

**Key words:** Bovine Leptospirosis, MAT, Serovars, Andaman.

### INTRODUCTION

Leptospirosis, a spirochaetal zoonosis, has emerged as a serious global veterinary and public health problem. The disease has gained much importance as it is often undiagnosed (Iqbal *et al.*, 2011). Bovine leptospirosis causes financial losses to the animals from the still birth, infertility, birth of weak calves, reduced milk yield and production (Grooms 2006). The first successful isolation of leptospira in India was reported among construction workers in the village of South Andaman by Taylor and Goyle (1931). However, the first identification of leptospira was done by Adinarayanan *et al.*

(1980) from Uttar Pradesh. In India, first isolation of leptospira by MAT was carried by Srivastava (1988) in man and animal from Andhra Pradesh.

Transmission of leptospirosis in human and animals occurred by the exposure to water or soil contaminated by the urine of infected animals or by direct contact with infected animals (Faine *et al.*, 1999). The genus *Leptospira* consists of both pathogenic and non-pathogenic species. The pathogenic leptospira is divided into more than 250 different serovars and the non-pathogenic into more than 60 serovars (Cerqueira and Picardeau 2009). Cattle

---

<sup>1</sup>Specialist Sero-epidemiology, Regional Disease Diagnostic Laboratory (ER), Kolkata, West Bengal, India.

<sup>2</sup>Assistant Director, ARD, Infectious Abortion Scheme Laboratory, IAH&VB, Kolkata, West Bengal, India.

<sup>3</sup> In charge, All India Co-ordinated Research Project on Foot and Mouth Disease, Kolkata Regional Centre, IAH&VB, Kolkata, West Bengal, India.

\*corresponding author. e - mail: drsumitvo@gmail.com.

are not only infected as accidental host but also maintenance host of specific *Leptospira* serovar strains and serves as reservoir animal.

## **MATERIALS AND METHODS**

### **Serum samples**

Serum samples of 108 cattle were sent from Senior Veterinary Officer (AH), Junglighat, Port Blair, Andaman and Nicobar Island to Regional Disease Diagnostic Laboratory (ER), Belgachia, Kolkata. The samples were collected in the months of January, 2014 and tested by Microscopic Agglutination Test (MAT) against eight serovars of *Leptospira interrogans*, namely *hardjo*, *pyrogenes*, *pomona*, *sejroe*, *bataviae*, *patoc*, *autumnalis*, and *australis*.

### **Leptospira media and culture**

Ellinghausen McCullough Johnson and Harris (EMJH) liquid media was prepared as per the standard protocol for the propagation of the leptospira culture. MAT was performed on all sera using eight reference serovars obtained from WHO National Reference Laboratory, Regional Medical Research Centre (RMRC), Port Blair, India.


### **Microscopic Agglutination Test:**

MAT was carried out as described by Cole *et al.* (1973) with some modification at Infectious Abortion Scheme (IAS) Laboratory, Institute of Animal Health & Veterinary Biologicals (IAH&VB), Kolkata. *Leptospira* live antigens required for MAT were grown in EMJH media in IAS Laboratory. For this, serovars were incubated at 30°C for 7 days in EMJH medium. Five to seven day-old cultures at a concentration of 1-2 x10<sup>8</sup> organisms /ml were used as live antigen for MAT. The concentration of *Leptospira* was measured in


Spectrophotometer. The sera were tested for the presence of specific antibodies against *Leptospira* serovars by MAT which was performed in 96 wells 'U' bottom "NUNC" Micro Titre plate. Firstly, all serum samples were initially diluted in three different dilutions 1:50, 1:100, 1:200 with phosphate buffer saline (PBS, pH7.4). After that, 100µl of each diluted serum was coated in to the U bottom micro titre plates, and then 100 µl of live antigen was added. The procedure was performed for all the eight serovars separately and incubated at 30°C for 2 hours. After that, a loopful of each mixture was examined under the dark field microscope (Leica) using 20X and 40X objectives for the presence of the agglutination. One antigen control and two (positive and negative) standard serum controls were used each time. Serum sample causing 50% agglutination at 1:100 dilutions or above was taken as positive reactor as per WHO/OIE manual for leptospirosis (Guitian *et al.*, 2001, Radostites *et al.*, 2007).

## **RESULTS AND DISCUSSION**


Among 108 sera samples 75 were positive to any one type of serovar by MAT as shown in Table 1. That finding also clearly indicated that overall 69.44% of serum samples were positive to one or more *Leptospira* species and 33 (31.56%) were negative. All the serum samples were tested in three dilutions (1:50, 1:100 & 1:200). The positive in agglutination shown as glistering woollen ball indicated the sera are very specific to *Leptospira* serovars as depicted in Fig.1 and Fig.2. The results of this study showed that most prevalent serovars were Autumnalis, Sejroe, Hardjo and Pomona (Table 2). The less prevalent serotypes are Pattoc, Bataviae, Pyrogenes and Australis (Table 2 and Fig.3). Among the serum samples tested, 34



**Fig. 1: MAT at 20X under Dark Field Microscope**




**Fig. 2: MAT at 40X under Dark Field Microscope.**



**Fig-3: Percentage of positive samples of *Leptospira* serovars.**

were positive to one serovar, 18 were positive to two serovars, 11 were positive to three serovars, 5 were positive to four serovars, 6 were positive to five serovars and 1 was positive to six serovars as shown in Table 1. These findings envisaged that most of the individual animals are infected with more than one serovar and this may cause great difficulties in controlling of the disease in that area (Fig.4).

Leptospirosis is known to occur in India since



**Fig-4: Incidence of one or more serovars of positive samples.**

early parts of 20<sup>th</sup> century. The disease is common in cattle and virtually in all the states of India (Srivastava and Kumar 2003). Many leptospirosis cases in animals and humans have been reported in the rainy season and after flood and heavy rainfall. Seasonal outbreak has also been reported in coastal areas especially in the Southern pennicular region, Maharashtra, 12.81% from Gujrat (Patel *et al.*, 2014), 42.50% from Odisha (Balamurgan *et al.*, 2013), 16.84%

**Table 1: Frequencies of one or more serovars of *Leptospira* sp.**

No of serovars	Numbers of positive serovars	Frequency (%)
One serovar	34	31.48
Two serovar	18	16.66
Three serovar	11	10.18
Four serovar	5	4.62
Five serovar	6	5.55
Six serovar	1	0.92
<b>Total</b>	<b>75</b>	<b>69.44</b>

from West Bengal (Mandal *et al.*, 2008). Knowledge of the prevalence of the *Leptospira* serovars is important for understanding the epidemiology of the disease and framing the public health policies aimed at prompt diagnosis and control measures. Studies of bovine leptospirosis in different parts of the world indicate that serovars responsible for reproductive losses vary depending on types of

serovars that are locally endemic since leptospiral antibodies may present in the serum for a considerable period of time after infection, the sero-reactivity may indicate the present or past exposure to leptospiral antigens (Balamurgan *et al.*, 2013).

MAT is considered as the gold standard test (Wolff, 1954) or International Test (Venkataraman *et al.*, 1992) for the diagnosis of leptospirosis. It is serovar specific test and choice for sero-epidemiological studies for detecting both IgG and IgM antibodies in animal sera. The sensitivity and specificity of MAT reported in a recent study were 91.94% and 73.77%, respectively (Dassanayake *et al.*, 2009).

It is well known fact that Hardjo serovar is common in cattle (Leonard *et al.*, 2004) and present study also correlates the similar findings. In addition to this, the prevalence of serovars Autumnalis (53.70%) was highest among the all serovars in bovine species. This finding corroborates with the findings of Sarvanan *et al.* (2000) who reported that

**Table 2: Frequency distribution of *Leptospira* serovars.**

Serovars	No. of samples reacted (1:100 dilution)	% positive samples against total samples	% frequencies against total no. of positive samples
Autumnalis	58	53.70	77.33
Sejroe	31	28.70	41.33
Hardjo	24	22.22	32.00
Pomona	23	21.29	30.66
Pattock	13	12.03	17.33
Bataviae	5	4.63	6.66
Pyrogenes	5	4.63	6.66
Australis	2	1.85	2.66

majority of human disease caused by *L. interrogans* serovar Autumnalis. In another study on incidence of *Leptospira interrogans* var *autumnalis* in canine was highest from West Bengal by Pal *et al.*(2013). However, the highest antibody titre against the three serovars namely Autumnalis, Sejroe and Hardjo indicate that the animal population was recently affected by these three serotypes. Although, direct transmission through skin abrasions occurs, it is evident that the contaminated environment or water bodies infected by the discharge of the urine or aborted materials by either infected animals or apparently healthy carrier animals plays an important role in disease spreading. Subsequently, the animals might have been exposed to the leptospira through the reservoir animals like rodents or other infected animals shedding the leptospira in the water bodies (Balamurgan *et al.*, 2013).

### CONCLUSION

High prevalence of bovine leptospirosis in Andaman Nicobar Island indicates its significance as endemic condition. Its prevalence in apparently healthy bovine indicates the presence of the agent in the environment which may be a potential zoonotic risk to human. This study also determines the need for continuous monitoring of leptospira burden in animals and human in close proximity to each other to combat the zoonotic infection.

### ACKNOWLEDGEMENTS

The authors are thankful to the Director, Animal Husbandry and Veterinary Services, Govt. of West Bengal and the Joint Director, ARD, Institute of Animal Health and Veterinary Biologicals for providing all the facilities to carry out this work.

### REFERENCES

- Adinarayanan N, James PC (1980) Studies on Leptospirosis in Farm stock and wild life in Kerala. *Ind Vet J* 57(6): 496-500.
- Balamurgan V, Thirumalesh SRA, Sridevi R, Mohandoss G, Govindaraj G, Hemadri D, Gajendragad MR, Rahman H (2013) Seroprevalence of bovine leptospirosis in Odhisa, India. *World J Vet Sci* 1: 1-17.
- Cerqueira GM, Picardeau M (2009) A century of Leptospira strain typing. *Infect Genet Evol* 9: 760-768.
- Cole JR, Sulzer CR, Pursell AR (1973) Improved Micro technique for the Leptospiral Microscopic Agglutination Test. *Applied Microbiol* 25: 976-980.
- Dassanayake DLB, Wimalaratna H, Agampodi SB, Liyanapathirana VC, Piyarathna TACL, Goonapienuwala BL (2009) Evaluation of surveillance case definition in the diagnosis of leptospirosis, using the microscopic agglutination test: a validation study. *BMC Infect Dis* 9: 48.
- Faine S, Adler B, Bolin C, Perolat P (1999) *Leptospira and Leptospirosis*. 2<sup>nd</sup> ed., Medi. Sci., Melbourne. Cited in Koizumi Nobuo, Yasutomi Ichiro (2012) Prevalance of leptospirosis in farm animals. *Japanese J Vet Res* 60(Suppl.): S55-S58.
- Grooms LD (2006) Reproductive losses caused by bovine viral diarrhoea virus and leptospirosis. *Theriogenology* 66: 624-628.
- Guitian FJ, Garcia-Pena FJ, Oliveira J, Sanjuan ML, Yus E (2001) Serological study of the frequency of leptospiral infections among dairy cows in farms with sub-optimal reproductive efficiency in Galicia, Spain. *Vet Microbiol* 80: 275-284.

Sero-prevalence of Bovine Leptospirosis in South Anandaman islands, India.

- Iqbal A, Singh R, Wazir VS, Ahmad I (2011) Current status of leptospirosis in dogs - an overview. *Ind Pet J* 3(1): 13-23.
- Leonard N, Mee JF, Snijders S, Mackie D (2004) Prevalence of antibodies to *Leptospira interrogans* serovar hardjo in bulk tank milk from unvaccinated Irish dairy herds. *Irish Vet J* 57: 226-231.
- Mandal S, Joardar SN, Chakraborty D, Sardar N (2008) Sero-epidemiological study of bovine leptospirosis in West Bengal. *Indian J Comp Microbiol Immunol Infect Dis* 29(1&2):42-44.
- Pal B, Mitra J, Roy BB, Basak D (2013). Seroprevalance of canine Leptospirosis in kennels of Kolkata. *Indian J Anim Hlth* 52(1): 27-30.
- Patel JM, Vihol PD, Prasad MC, Kalyani IH, Raval JK, Patel KM, Sushma RA, Thirumalesh, Balamurgan V (2014). Sero-epidemiological pattern of Leptospirosis in bovine of South Gujrat, India. *Vet World* 7(11): 999-1003.
- Radostites OM, Gay CC, HinChcliff KW, Constable PD (2007) *Veterinary Medicine* 10<sup>th</sup> Edn, Saunders Elsevier, Philadelphia, USA. 1286-1297.
- Sarvanan R, Rajendran P, Thyagarajan SP, Smythe LD, Norris MA, Simonds ML, Dohnt MF (2000) *Leptospira autumnalis* isolated from a human case from Avadi, India and the serovar's predominance in local rat and bandicoot populations. *Annual Tropical Med Parasitol* 94(5): 503-506.
- Srivastava SK (1988) Factors affecting anti-leptospira agglutinin titers in sera of animals and man. *Ind J Anim Sci* 58(6): 632-634.
- Srivastava SK, Kumar AA (2003) Seroprevalence of leptospirosis in animals and human beings in various regions in the country. *Ind J Comp Microbiol Immunol Infect Dis* 24: 155-159.
- Taylor J, Goyle AN (1931) Leptospirosis in Andamans. *Indian Medical Research Memoirs, Supplementary series to the Indian Journal of Medical Research. Ind J Med Res Memoir* 20: 55-56.
- Venkataraman KS, Nedunchelliyam S, Ramkrishna J, Ramdass P, Raghavan N (1992) Isolation of *Canicola leptospiral* serovar from urine of dog. *Ind Vet J* 69 (9): 866.
- Wolff JW (1954) *The laboratory diagnosis of leptospirosis*. C.C. Thomas, Springfield, Vol. III. Cited in Pal B, Mitra J, Roy BB, Basak D (2013) Sero prevalence of Canine leptospirosis in kennels of Kolkata. *Ind J Anim Health* 52(1):27-30.

**\*Cite this article as:** Mitra J, Chowdhury S, Pattanayak S (2015) Sero-prevalence of bovine Leptospirosis in South Anandaman islands, India. *Explor Anim Med Res* 5(1): 96-101.