

## Research Article

# THERAPEUTIC EFFECTS OF FEEDING GILOY (*TINOSPORA CORDIFOLIA*) AND FENUGREEK (*TRIGONELLA FOENUM-GRAECUM*) ON BLOOD BIOCHEMICAL PROFILE IN JERSEY CROSSBRED COWS

Anurag Sharma<sup>1\*</sup>, Naresh Kumar<sup>2</sup>, Ravindra Kumar<sup>3</sup>, Anika Sharma<sup>1</sup>

Received 07 February 2018, revised 02 May 2018

**ABSTRACT:** The study was conducted to evaluate the effect of Giloy (*Tinospora cordifolia*) and Fenugreek (*Trigonella foenum-graecum*) supplementation on blood biochemical parameters. The lactating cows were randomly divided into four groups, each group having six animals. T<sub>1</sub>, T<sub>2</sub> and T<sub>3</sub> group cows received Giloy stem powder (150 g), Fenugreek seed powder (150 g), and a combination of both the herbs (75 g of each) respectively, mixed with the concentrate feed for 60 days while the control group (T<sub>0</sub>) received only the concentrate. Blood sampling was done fortnightly, from Day 0 to Day 75. Blood plasma samples were analyzed for biochemical parameters viz. glucose, protein profile (Total protein, albumin, globulin and A: G ratio) and renal profile (urea nitrogen and creatinine). It was found that the feeding of Giloy stem powder lead to significantly lower levels ( $p \leq 0.05$ ) of blood glucose in the blood of lactating dairy cows. Most of the other blood biochemical parameters remained unchanged in control and treated animals.

**Key words:** Giloy (*Tinospora cordifolia*), Fenugreek (*Trigonella foenum-graecum*), Blood biochemical profile, Jersey crossbred.

## INTRODUCTION

The Himalayas are a rich repository of medicinal herbs. However, the areas with rich biodiversity remain potentially unexplored. The traditional knowledge about wild and cultivated veterinary medicinal plants used by indigenous people has not been much documented in the state. Discovery and development of newer drugs to replace ineffective or less effective drugs is a continuous phenomenon in modern medicine (Pattanayak *et al.* 2016). Giloy (*Tinospora cordifolia*), also known as *guduchi*, occupies the top spot in “*Ayurvedic Materia Medica*” and it has been designated as “*Rasayana*” (Bhattacharyya and Bhattacharya 2013). This plant finds mention in ancient Sanskrit literature like *Charak Samhita* and *Sushruta Samhita* as a potential healer of many diseases. The root extract of *T. cordifolia* is shown to exert hypolipidemic activity (Stanley *et al.* 1999), hypoglycaemic activity and to reduce serum and tissue cholesterol, phospholipids and free fatty acids (Stanley *et al.* 2000). The alcoholic extract of *T. cordifolia* (500mg/

kg body weight, orally) decreased the increased levels of serum creatinine, blood urea nitrogen and alkaline phosphatase in cisplatin induced nephrotoxicity in rats (Khanam *et al.* 2011). *T. cordifolia* supplementation in lactating Murrah buffaloes caused a higher plasma catalase activity and growth hormone concentration (Mir *et al.* 2013). *T. cordifolia* supplemented group in Murrah buffaloes showed that plasma glucose concentration was higher ( $p > 0.05$ ) than the control (Mir *et al.* 2014).

Fenugreek (*Trigonella foenum-graecum*) is a leguminous herb cultivated in India. The endosperm of the seed is rich in galactomannan and the young seeds mainly contain carbohydrates and sugar. Mature seeds contain the amino acid, fatty acid, vitamins and saponins. The main chemical constituents of fenugreek are fibres, flavonoids, polysaccharides, saponins and polysaccharides and some identified alkaloids such as trigonelline and choline (Toppo *et al.* 2009). Lower blood glucose ( $p < 0.05$ ) and urea ( $p < 0.01$ ) levels were reported in goats when fed on with fenugreek seeds (Alamer and

<sup>1</sup>Scholar <sup>2</sup>Professor (Retd), <sup>3</sup>Professor & Head, Department of Veterinary Physiology and Biochemistry, College of Veterinary and Animal Sciences, CSKHPKV, Palampur, Himachal Pradesh, India.

\*Corresponding author. e-mail: anurag.vets24@gmail.com

Basiouni 2005, Al-Janabi 2012). Fenugreek seeds have also been reported to lower the urea and creatinine content in plasma (Hamden *et al.* 2010). Blood plasma total protein, albumin, globulin, cholesterol, glucose and total lipids were found to have no significant difference on fenugreek supplementation in a study conducted on dairy goats (Al-Shaikh *et al.* 1999). Fenugreek seeds treatments showed significantly ( $p < 0.05$ ) increased blood glucose, total protein, albumin and creatinine in buffaloes (Abo El-Nor *et al.* 2007). Contrarily, fenugreek seeds have been evaluated for significantly decreased serum glucose, creatinine levels in streptozotocin-induced diabetic rats (Eidi *et al.* 2007). Treatment with fenugreek seed powder restored the levels of serum urea and creatinine levels as well as alkaline phosphatase, aspartate aminotransferase and alanine aminotransferase activities in gamma radiation irradiated rats which showed an increase in these biochemical parameters post exposure (El-Tawil 2009). The scientific literature lacks reports of effects of the herbs in dairy cattle and thus study was designed to evaluate their therapeutic effects in lactating dairy animals.

## MATERIALS AND METHODS

The study trial was conducted on Jersey crossbred, healthy cows in various stages of lactation, maintained at the Instructional Livestock Farm, College of Veterinary and Animal Sciences, CSKHPKV, Palampur (Himachal Pradesh), India after approval of study from Directorate of Research, CSKHPKV. Necessary permission for conducting the trial was obtained from In charge, Instructional Livestock Farm. The experimental animals were maintained in loose housing system, under standard feeding and management conditions being followed at Livestock farm. The animals were fed twice daily and watered *ad libitum*. The major fodder provided to the cows during entire study consisted of Setaria, Maize, Sorghum, local grass. In addition, the animals were also offered concentrate during milking time.

The Jersey crossbred lactating cows randomly divided into four groups, each group having six animals. T<sub>1</sub>, T<sub>2</sub> and T<sub>3</sub> group cows received Giloy stem powder (150 g), or Fenugreek seed powder (150 g), and a combination of both the herbs (75 g of each) respectively, mixed with the concentrate feed for 60 days while the control group (T<sub>0</sub>) received only the concentrate. The herbal treatment was administered at a fixed time daily *i.e.* afternoon milking hours, to all the animals till day 60 of the experimental trial. Blood samples from experimental dairy cows were collected at regular intervals of 15 days till Day 75 of the experiment. First sampling was done a day before the start (Day 0) of feeding trial. Sampling on Day 75 was done to evaluate residual effect of treatment, if any. The harvested blood plasma was then stored at -20°C till further analysis. The blood samples were taken by aseptic venipuncture technique (jugular vein) and were collected in heparinised tubes. Blood plasma biochemical parameters viz. Glucose, Protein profile (Total protein, Albumin, Globulin) and Renal profile (Urea Nitrogen, Creatinine) were determined using biochemical estimation kits (Manufactured by Agappe Diagnostics Ltd., Agappe Hills, Distt. Ernakulam Kerala, India) on automatic blood biochemistry analyzer (Mispa Nano, Agappe) on automatic blood biochemistry analyzer (Mispa Nano, Agappe). The results so obtained in the study were analyzed using computer software 'SAS Enterprise Guide'. The data was analyzed by using ANOVA at 5 % level of significance.

## RESULTS AND DISCUSSION


### Plasma glucose (mg/dL)

Glucose (mg/dL) in the blood plasma of lactating cows treated with herbal supplements and control group are depicted below in Fig. 1 and tabulated in Table 2.

Blood glucose concentration remained in normal physiological range (Kaneko *et al.* 2008). The control (T<sub>0</sub>) group did not show much variation in blood glucose during first 60 days. A significantly elevated blood

**Table 1. Composition of concentrate feed supplied to dairy cows (per 100kg).**

Ingredient	Quantity (kg)	Ingredient	Quantity (kg)
Maize	30	Mineral Mixture	3
Wheat Bran	10	Urea	1
Deoiled Rice Bran	15	Cottonseed Cake	8
Ground Nut Cake	10	Soya flakes	5
Lime Powder	1	Mustard Cake	9
Molasses	6	Bypass fat	1
Salt	1		
<b>Total</b>			<b>100</b>



**Fig. 1. Glucose (mg/dL) in the blood plasma of lactating cows treated with herbal supplements and control group.**

glucose level was observed on day 75. The blood plasma glucose concentration in giloy fed cows ( $T_1$  group) was significantly ( $p < 0.05$ ) lower as compared to control group from day 15 onwards.  $T_2$  and  $T_3$  groups did not exhibit any significant variation in blood glucose. An interesting observation was the significant increase in blood glucose values on day 75. As this elevation occurred in control as well treatment groups, it cannot be linked to herbal supplementation.

An increase, as well as a decline in blood glucose levels after herbal supplementation, has been reported by various research workers. Wadood *et al.* (1991) and Stanley *et al.* (2003) documented a decline in blood

glucose levels on *T. cordifolia* extract administration in lab animal studies. However, Mir *et al.* (2014) reported higher blood glucose levels in Murrah buffaloes in *T. cordifolia* treated group. Alamer and Basiouni (2005) found lower blood glucose in fenugreek fed goats. Contrarily, Abo El-Nor *et al.* (2007) reported an increase in blood glucose levels in fenugreek fed buffaloes. Vijayakumar *et al.* (2005) and Hamden *et al.* (2010) documented a decline in blood glucose levels on fenugreek extract administration in lab animal studies. Giloy (*Tinospora cordifolia*) is a documented herbal galactagogue (Mallick and Prakash 2011, Mir *et al.* 2014). Various studies reported a negative correlation between blood glucose and milk yield (Mouffok *et al.* 2013, Chládek and Máchal 2004, Djokoviæ *et al.* 2017). Thus the decline in the blood glucose level can be attributed to the enhanced glucose demand by mammary gland due to higher milk yield in the crossbred cows.

### Plasma protein profile

Total Protein, Albumin, Globulin and Albumin: Globulin ratio in the blood plasma of lactating cows treated with herbal supplements and control are tabulated in Table 3, 4, 5 and 6.

The values for total protein were slightly higher than standard reference (6.74 to 7.46 g/dL) for cattle (Kaneko

**Table 2. Glucose (mg/dL) in the blood plasma of lactating cows treated with herbal supplements and control group (Mean±S.E.).**

Group	Day 0	Day 15	Day 30	Day 45	Day 60	Day 75
$T_0$	54.03 <sup>bc</sup> ±0.62	51.38 <sup>cx</sup> ±1.79	53.55 <sup>bcx</sup> ±0.65	52.90 <sup>bcx</sup> ±1.02	55.95 <sup>bx</sup> ±1.97	65.57 <sup>ax</sup> ±0.69
$T_1$	56.22 <sup>a</sup> ±1.82	43.60 <sup>cy</sup> ±2.07	43.63 <sup>cy</sup> ±1.20	45.87 <sup>bcy</sup> ±0.78	49.10 <sup>by</sup> ±0.76	58.53 <sup>ay</sup> ±2.17
$T_2$	51.57 <sup>a</sup> ±2.17	48.22 <sup>axy</sup> ±1.96	48.75 <sup>axy</sup> ±2.85	50.48 <sup>axy</sup> ±2.76	49.93 <sup>ay</sup> ±2.65	55.15 <sup>ay</sup> ±2.42
$T_3$	52.77 <sup>b</sup> ±1.39	48.75 <sup>bxy</sup> ±2.04	48.92 <sup>bxy</sup> ±2.31	52.77 <sup>bx</sup> ±1.58	50.40 <sup>bxy</sup> ±1.69	65.30 <sup>ax</sup> ±1.61

Figures with different superscripts (a, b, c) differ significantly ( $p < 0.05$ ) between rows.

Figures with different superscripts (x, y, z) differ significantly ( $p < 0.05$ ) between columns.

$T_0$ -Control,  $T_1$ - Giloy,  $T_2$ -Fenugreek,  $T_3$ - Both (Fenugreek+Giloy).

**Table 3. Total Protein (g/dL) in the blood plasma of lactating cows treated with herbal supplements and control group (Mean±S.E.).**

Group	Day 0	Day 15	Day 30	Day 45	Day 60	Day 75
$T_0$	7.66 <sup>a</sup> ±0.19	7.40 <sup>a</sup> ±0.11	7.60 <sup>a</sup> ±0.09	7.55 <sup>a</sup> ±0.07	7.60 <sup>a</sup> ±0.09	7.60 <sup>a</sup> ±0.09
$T_1$	7.58 <sup>a</sup> ±0.17	7.17 <sup>a</sup> ±0.25	7.45 <sup>a</sup> ±0.16	7.50 <sup>a</sup> ±0.15	7.43 <sup>a</sup> ±0.19	7.45 <sup>a</sup> ±0.24
$T_2$	7.45 <sup>a</sup> ±0.11	7.10 <sup>a</sup> ±0.29	7.65 <sup>a</sup> ±0.19	7.35 <sup>a</sup> ±0.18	7.30 <sup>a</sup> ±0.13	7.33 <sup>a</sup> ±0.25
$T_3$	7.42 <sup>ab</sup> ±0.19	7.17 <sup>b</sup> ±0.18	7.60 <sup>a</sup> ±0.24	7.47 <sup>ab</sup> ±0.23	7.30 <sup>ab</sup> ±0.21	7.30 <sup>ab</sup> ±0.40

Figures with different superscripts (a, b, c) differ significantly ( $p < 0.05$ ) between rows.

Figures with different superscripts (x, y, z) differ significantly ( $p < 0.05$ ) between columns.

$T_0$ -Control,  $T_1$ - Giloy,  $T_2$ -Fenugreek,  $T_3$ - Both (Fenugreek+Giloy).

**Table 4. Albumin (g/dL) in the blood plasma of lactating cows treated with herbal supplements and control group (Mean±S.E.).**

Group	Day 0	Day 15	Day 30	Day 45	Day 60	Day 75
T <sub>0</sub>	3.23 <sup>a</sup> ±0.05	3.00 <sup>bxy</sup> ±0.06	3.10 <sup>bx</sup> ±0.03	3.13 <sup>abx</sup> ±0.02	3.08 <sup>b</sup> ±0.05	3.08 <sup>b</sup> ±0.03
T <sub>1</sub>	3.12 <sup>a</sup> ±0.05	2.87 <sup>by</sup> ±0.04	2.97 <sup>aby</sup> ±0.03	2.98 <sup>aby</sup> ±0.02	2.97 <sup>ab</sup> ±0.07	2.97 <sup>ab</sup> ±0.11
T <sub>2</sub>	3.17 <sup>a</sup> ±0.07	2.95 <sup>abxy</sup> ±0.03	3.10 <sup>abx</sup> ±0.06	3.13 <sup>abx</sup> ±0.05	2.87 <sup>b</sup> ±0.16	2.97 <sup>ab</sup> ±0.10
T <sub>3</sub>	3.27 <sup>a</sup> ±0.05	3.05 <sup>abx</sup> ±0.04	3.18 <sup>ax</sup> ±0.02	3.22 <sup>ax</sup> ±0.03	2.88 <sup>b</sup> ±0.10	3.10 <sup>a</sup> ±0.11

**Table 5. Globulin (g/dL) in the blood plasma of lactating cows treated with herbal supplements and control group (Mean±S.E.).**

Group	Day 0	Day 15	Day 30	Day 45	Day 60	Day 75
T <sub>0</sub>	4.43 <sup>a</sup> ±0.16	4.40 <sup>a</sup> ±0.11	4.50 <sup>a</sup> ±0.11	4.43 <sup>a</sup> ±0.10	4.53 <sup>a</sup> ±0.14	4.53 <sup>a</sup> ±0.14
T <sub>1</sub>	4.47 <sup>a</sup> ±0.16	4.30 <sup>a</sup> ±0.22	4.48 <sup>a</sup> ±0.16	4.52 <sup>a</sup> ±0.14	4.46 <sup>a</sup> ±0.16	4.48 <sup>a</sup> ±0.15
T <sub>2</sub>	4.28 <sup>a</sup> ±0.14	4.15 <sup>a</sup> ±0.27	4.55 <sup>a</sup> ±0.21	4.22 <sup>a</sup> ±0.20	4.43 <sup>a</sup> ±0.27	4.37 <sup>a</sup> ±0.20
T <sub>3</sub>	4.15 <sup>a</sup> ±0.21	4.12 <sup>a</sup> ±0.21	4.42 <sup>a</sup> ±0.22	4.25 <sup>a</sup> ±0.21	4.42 <sup>a</sup> ±0.22	4.20 <sup>a</sup> ±0.32

**Table 6. Albumin: Globulin ratio in the blood plasma of lactating cows treated with herbal supplements and control group (Mean±S.E.).**

Group	Day 0	Day 15	Day 30	Day 45	Day 60	Day 75
T <sub>0</sub>	0.73 <sup>a</sup> ±0.03	0.68 <sup>a</sup> ±0.02	0.69 <sup>a</sup> ±0.02	0.71 <sup>axy</sup> ±0.03	0.68 <sup>a</sup> ±0.02	0.68 <sup>a</sup> ±0.02
T <sub>1</sub>	0.70 <sup>a</sup> ±0.03	0.67 <sup>a</sup> ±0.03	0.67 <sup>a</sup> ±0.03	0.66 <sup>ay</sup> ±0.02	0.67 <sup>a</sup> ±0.01	0.66 <sup>a</sup> ±0.02
T <sub>2</sub>	0.75 <sup>a</sup> ±0.04	0.73 <sup>a</sup> ±0.05	0.69 <sup>a</sup> ±0.04	0.75 <sup>axy</sup> ±0.04	0.65 <sup>a</sup> ±0.07	0.68 <sup>a</sup> ±0.03
T <sub>3</sub>	0.80 <sup>a</sup> ±0.05	0.75 <sup>a</sup> ±0.04	0.73 <sup>a</sup> ±0.04	0.77 <sup>ax</sup> ±0.04	0.65 <sup>a</sup> ±0.06	0.75 <sup>a</sup> ±0.05

**Table 7. Urea Nitrogen (mg/dL) in the blood plasma of lactating cows treated with herbal supplements and control group (Mean±S.E.).**

Group	Day 0	Day 15	Day 30	Day 45	Day 60	Day 75
T <sub>0</sub>	23.98 <sup>by</sup> ±2.14	27.28 <sup>by</sup> ±1.02	29.65 <sup>ab</sup> ±2.41	34.23 <sup>a</sup> ±1.72	28.48 <sup>abx</sup> ±0.74	29.78 <sup>ab</sup> ±2.47
T <sub>1</sub>	29.83 <sup>ax</sup> ±1.00	32.83 <sup>ax</sup> ±2.16	32.02 <sup>a</sup> ±2.07	32.03 <sup>a</sup> ±3.56	26.03 <sup>axy</sup> ±1.44	30.52 <sup>a</sup> ±3.23
T <sub>2</sub>	25.95 <sup>bxy</sup> ±1.20	27.57 <sup>by</sup> ±0.85	28.45 <sup>b</sup> ±0.78	34.10 <sup>a</sup> ±1.99	24.33 <sup>by</sup> ±0.83	27.60 <sup>b</sup> ±2.76
T <sub>3</sub>	26.18 <sup>bxy</sup> ±1.50	29.25 <sup>abxy</sup> ±1.29	32.25 <sup>a</sup> ±2.16	34.05 <sup>a</sup> ±1.64	24.58 <sup>cy</sup> ±1.49	31.80 <sup>ab</sup> ±2.93

**Table 8. Creatinine (mg/dL) in the blood plasma of lactating cows treated with herbal supplements and control group (Mean±S.E.).**

Group	Day 0	Day 15	Day 30	Day 45	Day 60	Day 75
T <sub>0</sub>	0.75 <sup>a</sup> ±0.05	0.78 <sup>a</sup> ±0.05	0.75 <sup>a</sup> ±0.04	0.78 <sup>a</sup> ±0.03	0.83 <sup>a</sup> ±0.06	0.83 <sup>a</sup> ±0.05
T <sub>1</sub>	0.72 <sup>a</sup> ±0.07	0.67 <sup>a</sup> ±0.04	0.72 <sup>a</sup> ±0.03	0.73 <sup>a</sup> ±0.02	0.77 <sup>a</sup> ±0.04	0.82 <sup>a</sup> ±0.07
T <sub>2</sub>	0.75 <sup>a</sup> ±0.04	0.67 <sup>a</sup> ±0.03	0.73 <sup>a</sup> ±0.02	0.72 <sup>a</sup> ±0.02	0.78 <sup>a</sup> ±0.05	0.77 <sup>a</sup> ±0.12
T <sub>3</sub>	0.79 <sup>ab</sup> ±0.03	0.73 <sup>b</sup> ±0.02	0.83 <sup>ab</sup> ±0.06	0.77 <sup>ab</sup> ±0.02	0.78 <sup>ab</sup> ±0.05	0.92 <sup>a</sup> ±0.07

In Table No. 4, 5, 6, 7, 8, figures with different superscripts (a, b, c) differ significantly ( $p < 0.05$ ) between rows.

Figures with different superscripts (x, y, z) differ significantly ( $p < 0.05$ ) between columns.

T<sub>0</sub>-Control, T<sub>1</sub>- Giloy, T<sub>2</sub>-Fenugreek, T<sub>3</sub>-Both (Fenugreek+Giloy).

*et al.* 2008). The plasma total protein values in control ( $T_0$ ) and treatment groups ( $T_1$ ,  $T_2$  and  $T_3$ ) were found to be in normal range and did not show much variation during the trial.

The albumin values are in agreement with the standard reference range for cattle (3.03 to 3.55 g/dL) given by Kaneko *et al.* (2008). The plasma albumin values in the case of control ( $T_0$ ) group and treatment groups ( $T_1$ ,  $T_2$  and  $T_3$ ) the albumin values declined on day 15 to a lower value and remained low within the group for entire study period with no particular trend observed. The albumin values from day 15 to day 45 in the  $T_1$  group were found to be lower as compared to control group. However, in  $T_2$  and  $T_3$  albumin values were similar to the corresponding values of the control group. The albumin levels did not vary much after thereafter.

The plasma globulin (g/dL) in the blood plasma is higher than the standard reference values (3.00 to 3.48 g/dL) for cattle (Kaneko *et al.* 2008). Frequent vaccination and exposure to various microbes (in comparison to the animals for which the reference values are quoted) might be the reason for higher globulin values in Jersey crossbred cows used in the present study. The variations observed were statistically non-significant.

In general the A: G ratio declined slightly from day 15 onwards in all the groups. However, the differences were non-significant. The A: G value is indicative of the immune status of the animal. The decline in A: G values is attributable to increase in plasma globulins which could be due to the fact that animals were vaccinated early at the beginning of the trial (day 5).

Maher and NMB (2013) and Al-Shaikh *et al.* (1999) reported no significant effect of fenugreek supplementation on plasma total protein levels in dairy cows and goats respectively. However, Nasser (2013) and Abo El-Nor *et al.* (2007) reported an increase in total protein levels on fenugreek supplementation in cows. Aher and Wahi (2010) observed that the immunomodulatory action of *T. cordifolia* lead to an increase in the immunoglobulin levels in rats. Similarly, it has been documented that fenugreek feeding in cows lead to elevated globulin levels (Nasser 2013). However, Maher and NMB (2013) found no significant difference in plasma globulin levels in Friesian cows after fenugreek supplementation.

### Plasma renal profile

The plasma urea nitrogen and creatinine levels (mg/dL) in the blood plasma of lactating cows treated with herbal supplements and control are tabulated in Table 7 and Table 8 respectively.

The standard reference range for plasma urea nitrogen is 20-30 mg/dL (Kaneko *et al.* 2008). Slightly higher values observed in the present study may be due to variation in dietary protein intake. No specific trend in the variation of plasma urea nitrogen and creatinine values could be seen in control ( $T_0$ ) as well as treatment ( $T_1$ ,  $T_2$  and  $T_3$ ) groups. The plasma urea nitrogen concentration in the cows receiving fenugreek seed powder or a combination of fenugreek and giloy ( $T_2$  and  $T_3$  groups) was found to be significantly lower on day 60. A decline in blood urea levels after herbal supplementation has been reported by various research workers. Nasser (2013) reported a decline in blood urea levels on fenugreek supplementation in the dairy cows. Alamer and Basiouni (2005) and Al-Janabi (2012) reported a decline in blood urea levels in goats fed with fenugreek seeds.

Giloy (*Tinospora cordifolia*) is a documented herbal galactagogue (Mallick and Prakash 2011, Mir *et al.* 2014). Also there is an evidence that there is a negative correlation between blood glucose and milk yield (Mouffok *et al.* 2013, Chládek and Máchal 2004, Djokoviæ *et al.* 2017). Thus the decline in the blood glucose level can be attributed to the enhanced glucose demand by mammary gland due to higher milk yield in the crossbred cows.

### CONCLUSION

The study can be concluded that the blood biochemical parameters showed some variation in some groups. In the giloy fed group ( $T_1$ ), the plasma glucose levels were significantly lower than the control group animals. The lactating dairy cows supplemented with fenugreek ( $T_2$ ) or both giloy and fenugreek ( $T_3$ ) did not exhibit any significant variation in plasma glucose levels during the supplementation period. Protein profile (total protein, albumin, globulin and A: G ratio) was not affected much during the course of giloy supplementation. However, lower levels of plasma albumin and higher levels of plasma globulins were seen from day 15 to day 45. A: G ratio was significantly lower in giloy fed animals in comparison to control on day 45. No significant variations were observed in the protein profile values of group  $T_2$  and  $T_3$  animals. No significant effect on the renal profile (urea nitrogen and creatinine) could be attributed to the giloy supplementation. Similarly, in animals receiving fenugreek or both the herbs ( $T_2$  and  $T_3$ ) urea nitrogen levels exhibited nonspecific variation which appears to be not linked to herbal supplementation.



## REFERENCES

- Abo El-Nor SAH, Khattab HM, Al-Alamy HA, Salem FA, Abdou MM (2007) Effect of some medicinal plants seeds in the rations on the productive performance of lactating buffaloes. Intern J Dairy Sci 2(4): 348-355.
- Aher VD, Wahi AK (2010) Pharmacological study of *Tinospora cordifolia* as an immunomodulator. Intern J Current Pharmaceut Res 2(4): 52-54.
- Alamer MA, Basiouni GF (2005) Feeding effects of fenugreek seeds (*Trigonella foenum-graecum*) on lactation performance, some plasma constituents and growth hormone level in goats. Pakistan J Biologic sci 8(11): 1553-1556.
- Al-Janabi, Abdul K, Ahmed F (2012) Feeding effects of fenugreek seeds (*Trigonella foenum-graecum*) on lactation performance, some serum constituents and prolactin hormone level in Damascus crossbred goats. Diyala Agric Sci J 4(1): 1-8.
- Al-Shaikh MA, Al-Mufarrej SI, Mogawer HH (1999) Effect of fenugreek seeds (*Trigonella foenum-graecum*) on lactational performance of dairy goat. J Applied Anim Res 16: 177-183.
- Bhattacharyya C, Bhattacharyya G (2013) Therapeutic potential of Giloy, *Tinospora cordifolia* (Willd.) Hook. f. & Thomson (Menispermaceae): The magical herb of Ayurveda. Intern J Pharmaceut Biologic Archiv 4(4): 558-584.
- Chládek G, Máchal L (2004) Changes in the relationship between blood plasma glucose concentration and milk production in Czech Pied cows in the course of the year. Acta Universitatis Agriculturae et Silviculturae Mendelianae Brunensis 2: 97-104.
- Djokovic R, Kuraubic V, Ilić Z, Cincovic, Lalovic M, Jašovic B, Bojkovski J (2017) Correlation between blood biochemical metabolites, milk yield, dry matter intake and energy balance in dairy cows during early and mid lactation. Advances Diabetes Metabolism 5(2): 26-30.
- Eidi A, Eidi M, Sokhteh M (2007) Effect of fenugreek (*Trigonella foenum-graecum*) seeds on serum parameters in normal and streptozotocin-induced diabetic rats. Nutrition Res 27: 728-733.
- El-Tawil GA (2009) Effect of fenugreek (*Trigonella foenum-graecum*) supplementation on radiation-induced oxidative stress in liver and kidney of rats. J Radiation Res Applied Sci 2(1): 19-30.
- Hamden K, Jaouadi B, Carreau S, Bejar S, Abdelfattah E (2010) Inhibitory effect of fenugreek galactomannan on digestive enzymes related to diabetes, hyperlipidemia, and liver-kidney dysfunctions. Biotechnol Bioprocess Enginee 15: 407-413.
- Kaneko J, Harvey J, Bruss M (2008) Clinical Biochemistry of Domestic Animals (6<sup>th</sup> edn.). Academic Press, Inc. 882-888.
- Khanam S, Mohan NP, Devi K, Sultana R (2011) Protective role of *Tinospora Cordifolia* against cisplatin induced nephrotoxicity. Interna J Pharmacy Pharmaceut Sci 3(4): 98-70.
- Maher A, NMB (2013) Effect of using Fenugreek seeds on milk yield and some blood characteristics in local friesian cows. J Tikrit Univ Agril Sci 13(2): 70-75.
- Mallick S, Prakash BS (2011) Influence of feeding *Tinospora cordifolia* peripartum on lactation parameters in crossbred cows. J Anim Physiol Anim Nutri 96(6): 1112-1120.
- Mir AN, Parveen K, Aarif O, Shergojry SA (2013) Effect of *Tinospora cordifolia* supplementation on certain biochemical parameters in lactating Murrah buffaloes during winter season. Intern J Scientif Res 2(7): 515-516.
- Mir AN, Parveen K, Wani SA, Shergojry SA, Ashutosh, Aarif O (2014) Milk production status of lactating murrah buffaloes on *Tinospora cordifolia* supplemented diet with special reference to immunological, metabolic and hormonal profile. Anim Sci Reporter 8(1): 18-25.
- Mouffok CE, Madani T, Semara L, Ayache N, Rahal A (2013) Correlation between body condition score, blood biochemical metabolites, milk yield and quality in Algerian Montbéliarde cattle. Pakistan Vet J 33(2): 191-194.
- Nasser AK, Shams Al-dain QZ, Abou, Mahmood (2013) Using fenugreek seeds powder as a feed additive in rations of Sharabi local cows and its effect on some hematological and biochemical parameters. Iraqi J Vet Sci 27(1): 13-19.
- Pattanayak S, Mandal TK, Bandyopadhyay SK (2016) Validation and therapeutic use of succulent plant parts - opening of a new horizon of alternative medicine. Explor Anim Med Res 6(1): 8-14.
- Stanely P, Prince M, Menon VP (2000) Hypoglycaemic and other related actions of *Tinospora cordifolia* roots in alloxan-induced diabetic rats. J Ethnopharmacology 70: 9-15.
- Stanely P, Prince M, Menon V P (2003) Hypoglycaemic and hypolipidaemic action of alcohol extract of *Tinospora cordifolia* roots in chemical induced diabetes in rats. Phytoterpay Res 17: 410-413.

Stanely P, Prince M, Menon VP, Gunasekaran PG (1999) Hypolipidaemic action of *Tinospora cordifolia* roots in alloxan diabetic rats. J Ethnopharmacol 64: 53-57.

Toppo FA, Akhand R, Pathak AK (2009) Pharmacological actions and potential uses of *Trigonella foenum-graecum*: A review. Asian J Pharmaceut Clinic Res 2(4): 29-38.

Vijayakumar MV, Singh S, Chhipa RR, Bhat MK (2005) The hypoglycaemic activity of fenugreek seed extract is mediated through the stimulation of an insulin signalling pathway. British J Pharmacol 146: 41-48.

Wadood N, Wadood A, Shah SAW (1991) Effect of *Tinospora cordifolia* on blood glucose and total lipid levels of normal and alloxan-diabetic rabbits. Planta medica 58: 131-136.

**\*Cite this article as:** Sharma A, Kumar N, Kumar R, Sharma A (2018) Therapeutic effects of feeding Giloy (*Tinospora cordifolia*) and Fenugreek (*Trigonella foenum-graecum*) on blood biochemical profile in Jersey crossbred cows. Explor Anim Med Res 8(1): 52-58.